

Acta Crystallographica Section E

## **Structure Reports Online**

ISSN 1600-5368

S. Mondal,<sup>a</sup> R. S. Narasegowda,<sup>b</sup> B. Nagaraj,<sup>b</sup> H. S. Yathirajan,<sup>b</sup> T. Narasimhamurthy<sup>c</sup> and Ravindranath S. Rathore<sup>d\*</sup>

<sup>a</sup>Department of Physics, India Institute of Science, Bangalore, India, <sup>b</sup>Department of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore 570 006, India, <sup>c</sup>Bioinformatics Center, Indian Institute of Science, Bangalore 560 012, India, and <sup>d</sup>Oriental Organization of Molecular and Structural Biology, 204 Agarwal Bhavan, Malleshwaram, Bangalore 560 055, India

Correspondence e-mail: ravindranath\_rathore@yahoo.com

#### **Key indicators**

Single-crystal X-ray study T = 293 KMean  $\sigma(\text{C-C}) = 0.003 \text{ Å}$  R factor = 0.051 wR factor = 0.150Data-to-parameter ratio = 17.9

For details of how these key indicators were automatically derived from the article, see http://journals.iucr.org/e.

# 2,2'-Dipropylmalonic acid: intermolecular association *via* strong O—H···O hydrogen bonds

The molecular structure of the title compound,  $C_9H_{16}O_4$ , is described by a *trans*-planar configuration of the propyl chains. The molecule has an approximate pseudo-twofold rotation axis in the crystal structure. The molecules are interconnected by strong  $O-H\cdots O$  hydrogen bonds and form an  $R_2^2(8)$  ring structure.

Received 6 December 2004 Accepted 13 December 2004 Online 24 December 2004


#### Comment

The molecular assembly in the title compound, (I), is predominantly governed by N-H···O, O-H···O and O-H···N hydrogen bonds (Desiraju, 2002). Quite often, hydrogen-bonded patterns are formed which are interesting in supramolecular design and crystal engineering. Such patterns of hydrogen bonds were first analysed by Etter using graph theory (Etter, 1990) and later the formalism was further developed by Bernstein *et al.* (1995). In this report, we discuss 2,2'-dipropylmalonic acid, which is an intermediate for the preparation of the anticonvulsant drug valproic acid (Abbott & Acheampong, 1988). The intermolecular linkage *via* O-H···O hydrogen bonds forms a ring structure.

The molecular structure of (I) is shown in Fig. 1. The molecular conformation is characterized by four intersecting planes formed by propyl chains and carboxylic acid groups, and the molecule adopts a sort of propeller-shape. The maximum out-of-plane deviations from the least-square planes formed by the non-H atoms in the propyl chains and carboxylic acid groups are within 0.03 Å. The conformation of (I) is described by the torsion angles  $C1-C2-C4-C5=-62.1\ (2)^\circ$ ,  $C3-C2-C7-C8=-56.9\ (2)^\circ$ ,  $C2-C4-C5-C6=179.0\ (2)^\circ$  and  $C2-C7-C8-C9=-177.9\ (2)^\circ$ . The last two are *trans*, indicating fully extended propyl chains. Compound (I) possesses molecular  $C_2$  point group symmetry. However, the molecular symmetry is only approximately retained in the crystal structure. The r.m.s. deviation between related components is  $0.06\ \text{Å}$ .

The hydrogen-bond parameters are given in Table 1 and a packing diagram is shown in Fig. 2. The molecules are interconnected by strong  $O-H\cdots O$  hydrogen bonds. Intermolecular interactions  $via\ O-H\cdots O$  bonds form an  $R_2^2(8)$  ring structure (Bernstein *et al.*, 1995).

© 2005 International Union of Crystallography Printed in Great Britain – all rights reserved



**Figure 1**The molecular structure of (I). Displacement ellipsoids are drawn at the 30% probability level and H atoms are shown as small circles of arbitrary radii




Figure 2 Packing diagram of (I), illustrating the intermolecular associations via strong  $O-H\cdots O$  hydrogen bonds (dashed lines), forming a hydrogen-bonded  $R_2^2(8)$  pattern. H atoms have been omitted.

#### **Experimental**

The title compound was synthesized by treating diethyl dipropylmalonate with sodium hydroxide at 353–363 K for 6 h. The melting point of (I) was 432 K. The compound was crystallized by slow evaporation of a saturated chloroform solution.

#### Crystal data

$C_9H_{16}O_4$	Z = 2
$M_r = 188.22$	$D_x = 1.145 \text{ Mg m}^{-3}$
Triclinic, $P\overline{1}$	Mo $K\alpha$ radiation
a = 7.683 (1)  Å	Cell parameters from 3369
b = 8.549 (1)  Å	reflections
c = 9.382 (1)  Å	$\theta = 5-55^{\circ}$
$\alpha = 76.249 (2)^{\circ}$	$\mu = 0.09 \text{ mm}^{-1}$
$\beta = 70.140 (2)^{\circ}$	T = 293 (2)  K
$\gamma = 72.355 (2)^{\circ}$	Block, colourless
$V = 546.05 (12) \text{ Å}^3$	$0.55 \times 0.22 \times 0.12 \text{ mm}$

#### Data collection

Bruker SMART CCD area-detector diffractometer	2261 independent reflections 1784 reflections with $I > 2\sigma(I)$
$\omega$ scans	$R_{\rm int} = 0.014$
Absorption correction: multi-scan	$\theta_{\rm max} = 27.4^{\circ}$
(SADABS; Sheldrick, 1996)	$h = -9 \rightarrow 9$
$T_{\min} = 0.977, T_{\max} = 0.989$	$k = -10 \rightarrow 11$
5840 measured reflections	$l = -12 \rightarrow 12$

#### Refinement

refinement

,	
Refinement on $F^2$	$w = 1/[\sigma^2(F_o^2) + (0.0805P)^2]$
$R[F^2 > 2\sigma(F^2)] = 0.051$	+ 0.0901P]
$wR(F^2) = 0.150$	where $P = (F_o^2 + 2F_c^2)/3$
S = 1.04	$(\Delta/\sigma)_{\rm max} < 0.001$
2261 reflections	$\Delta \rho_{\text{max}} = 0.18 \text{ e Å}^{-3}$
126 parameters	$\Delta \rho_{\min} = -0.16 \text{ e Å}^{-3}$
H atoms treated by a mixture of	
independent and constrained	

**Table 1** Hydrogen-bonding geometry (Å, °).

$D-H\cdots A$	D-H	$H \cdot \cdot \cdot A$	$D \cdot \cdot \cdot A$	$D-H\cdots A$
$O1-H1\cdots O3^{i}$	0.83 (3)	1.85 (3)	2.678 (2)	176 (3)
$O2-H2\cdots O4^{ii}$	0.87 (3)	1.80 (3)	2.673 (2)	175 (3)

Symmetry codes: (i) -x, -y, 2-z; (ii) -x, 1-y, 1-z.

The methyl and methylene H atoms were refined with fixed geometry (C-H = 0.96 and 0.97 Å), riding on their carrier atoms, with  $U_{\rm iso}$  values set at 1.2 (1.5 for the methyl H atoms) times  $U_{\rm eq}$  of the parent atom. The carboxy H atoms were located in a difference electron-density map and were refined isotropically.

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT-Plus* (Bruker, 2001); data reduction: *SAINT-Plus*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *PLUTON* (Spek, 1999); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PARST* (Nardelli, 1995).

We thank Professor T. N. Guru Row for providing access to the CCD facility.

#### References

 Abbott, F. S. & Acheampong, A. A. (1988). Neuropharmacology, 27, 287–94.
Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). Angew. Chem. Int. Ed. Engl. 34, 1555–1573.

Bruker (1998). SMART. Version 5.054. Bruker AXS Inc., Madison, Wisconsin, USA.

Bruker (2001). SAINT-Plus. Version 6.22. Bruker AXS Inc., Madison, Wisconsin, USA.

### organic papers

Desiraju, G. R. (2002). Acc. Chem. Res. 35, 565–573. Etter, M. C. (1990). Acc. Chem. Res. 23, 120–126. Farrugia, L. J. (1997). J. Appl. Cryst. 30, 565. Farrugia, L. J. (1999). J. Appl. Cryst. 32, 837–838. Nardelli, M. (1995). J. Appl. Cryst. 28, 659.

Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.

Sheldrick, G. M. (1997). SHELXL97 and SHELXS97. University of Göttingen, Germany.

Spek, A. L. (1999). PLUTON. University of Utrecht. The Netherlands.

Acta Cryst. (2005). E61, o151–o153 S. Mondal et al. •  $C_9H_{16}O_4$  **o153**