

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

6-(4-Fluorophenyl)-3-phenyl-7H-1,2,4-triazolo[3,4-b][1,3,4]thiadiazine

B. S. Palakshamurthy,^{a*} H. C. Devarajegowda,^a
N. R. Mohan,^b S. Sreenivasa^b and P. A. Suchetan^c^aDepartment of Physics, Yuvaraja's College (Constituent College), University of Mysore Mysore, Karnataka, India, ^bDepartment of Studies and Research in Chemistry, Tumkur University, Tumkur, Karnataka 572 103, India, and ^cDepartment of Studies and Research in Chemistry, U.C.S., Tumkur University, Tumkur, Karnataka 572 103, India

Correspondence e-mail: palaksha.bspm@gmail.com


Received 11 February 2014; accepted 13 February 2014

Key indicators: single-crystal X-ray study; $T = 294$ K; mean $\sigma(C-C) = 0.004$ Å; R factor = 0.053; wR factor = 0.134; data-to-parameter ratio = 10.8.

In the title compound, $C_{16}H_{11}FN_4S$, the dihedral angles between the triazole ring and the phenyl and fluorobenzene rings are 23.22 (17) and 18.06 (17)°, respectively. The six-membered heterocyclic ring adopts a distorted envelope conformation, with the methylene C atom as the flap. In the crystal, the molecules are linked by two $C-H \cdots N$ and $C-H \cdots F$ interactions along [010], forming $C(5)$, $C(8)$ and $C(13)$ chains respectively. $C-H \cdots \pi$ interactions involving the phenyl ring and $\pi-\pi$ interactions [centroid-centroid separation for triazole rings = 3.5660 (18) Å] are also observed.

Related literature

For the antifungal activity of nitrogen-containing heterocycles, see: Mathew *et al.* (2007) and for their antibacterial activity, see: Demirbas *et al.* (2005).



Experimental

Crystal data

$C_{16}H_{11}FN_4S$
 $M_r = 310.35$
 Monoclinic, $P2_1/c$
 $a = 15.088$ (2) Å
 $b = 13.464$ (2) Å
 $c = 7.0557$ (12) Å
 $\beta = 91.076$ (3)°
 $V = 1433.0$ (4) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.24$ mm⁻¹
 $T = 294$ K
 $0.27 \times 0.23 \times 0.18$ mm

Data collection

Bruker APEXII CCD diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 2007)
 $T_{min} = 0.939$, $T_{max} = 0.958$
 9962 measured reflections
 2154 independent reflections
 1630 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.060$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.134$
 $S = 0.94$
 2154 reflections
 199 parameters
 H-atom parameters constrained
 $\Delta\rho_{max} = 0.19$ e Å⁻³
 $\Delta\rho_{min} = -0.22$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg3 is the centroid of the C1–C6 phenyl ring.

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$C9-H9B \cdots N2^i$	0.97	2.36	3.301 (3)	164
$C12-H12 \cdots N2^i$	0.93	2.47	3.393 (4)	173
$C4-H4 \cdots F1^{ii}$	0.93	2.57	3.475 (4)	164
$C1-H1 \cdots Cg3^{iii}$	0.93	2.93	3.598 (3)	130

Symmetry codes: (i) $-x + 2, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x, y - 1, z$; (iii) $x, -y + \frac{3}{2}, z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2009); cell refinement: APEX2 and SAINT-Plus (Bruker, 2009); data reduction: SAINT-Plus and XPREP (Bruker, 2009); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 2012) and Mercury (Macrae *et al.*, 2008); software used to prepare material for publication: SHELXL97.

The authors thank H. T. Srinivasa, Raman Research Institute, Bangalore, India for useful discussions.

Supporting information for this paper is available from the IUCr electronic archives (Reference: HB7198).

References

- Bruker (2009). APEX2, SADABS, SAINT-Plus and XPREP. Bruker AXS Inc., Madison, Wisconsin, USA.
 Demirbas, N., Demibras, A., Karaoglu, S. A. & Celik, E. (2005). *Arkivoc*, **1**, 75–91.
 Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
 Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
 Mathew, V., Keshavayya, J., Vaidya, V. P. & Giles, D. (2007). *Eur. J. Med. Chem.* **42**, 823–840.
 Sheldrick, G. M. (2006). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2014). E70, o375 [doi:10.1107/S1600536814003262]

6-(4-Fluorophenyl)-3-phenyl-7H-1,2,4-triazolo[3,4-*b*][1,3,4]thiadiazine

B. S. Palakshamurthy, H. C. Devarajegowda, N. R. Mohan, S. Sreenivasa and P. A. Suchetan

S1. Comment

Nitrogen containing heterocyclic molecules show a broad spectrum of pharmacological properties like antifungal (Mathew *et al.*, 2007), antibacterial (Demirbas *et al.*, 2005) activities. As part of our studies in this area, the title compound was synthesized and its crystal structure determined.

In the title compound, C₁₄H₁₁FN₄S, the dihedral angle between the ring pairs A—B, A—C, A—D, B—C, B—D and C—D are 23.22 (16)°, 16.62 (13)°, 29.83 (16)°, 9.86 (14)°, 18.06 (16)° and 14.61 (14)° respectively. In the crystal, the molecules are linked into one another through C9—H9···N2, C12—H12···N2 and C4—H4···F1 interactions along [010] forming C(5), C(8) and C(13) chains respectively. The structure is further stabilized by C—H··· π and π ··· π interactions [centroid-centroid separation = 3.5660 Å] along [001] leading to a two dimensional architecture.

S2. Experimental

An equimolar mixture of 4-amino-5-phenyl-4*H*-1,2,4-triazole-3-thiol (1 mmole) and 2-chloro-1-(4-fluorophenyl)-ethanone (1 mmole) and sodium acetate (2.5 mmol) in absolute ethanol (10 ml) were refluxed for 2 h and completion of the reaction was monitored by TLC. Reaction mixture was cooled to room temperature, the solvent was removed under vacuum and the precipitate obtained was filtered, washed with water and dried to get crude product. The Crude solid was further purified by column chromatography using dichloromethane/methanol (9:1) as eluent and was later recrystallized from dichloromethane/methanol solvent system to get colorless prisms of the title compound.

S3. Refinement

The other H atoms were positioned with idealized geometry using a riding model with C—H = 0.93–0.96 Å. All H atoms were refined with isotropic displacement parameters (set to 1.2–1.5 times of the U eq of the parent atom).




Figure 1

Molecular structure of the title compound, showing displacement ellipsoids drawn at the 50% probability level.




Figure 2

Formation of C(5), C(8) and C(13) chains through C—H...N and C—H...F hydrogen bonds.




Figure 3
Display of C—H \cdots π and $\pi\cdots\pi$ interactions in the crystal structure.

6-(4-Fluorophenyl)-3-phenyl-7H-1,2,4-triazolo[3,4-*b*][1,3,4]thiadiazine

Crystal data

$C_{16}H_{11}FN_4S$

$M_r = 310.35$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 15.088$ (2) Å

$b = 13.464$ (2) Å

$c = 7.0557$ (12) Å

$\beta = 91.076$ (3)°

$V = 1433.0$ (4) Å³

$Z = 4$

$F(000) = 640$

Prism

$D_x = 1.438$ Mg m⁻³

Melting point: 523 K

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 199 reflections

$\theta = 1.2$ – 26°

$\mu = 0.24$ mm⁻¹

$T = 294$ K

Prism, colourless

$0.27 \times 0.23 \times 0.18$ mm

Data collection

Bruker APEXII CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
Detector resolution: 1.6 pixels mm⁻¹
phi and ω scans
Absorption correction: multi-scan
(*SADABS*; Sheldrick, 2007)
 $T_{\min} = 0.939$, $T_{\max} = 0.958$

9962 measured reflections
2154 independent reflections
1630 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.060$
 $\theta_{\max} = 26.0^\circ$, $\theta_{\min} = 2.0^\circ$
 $h = -18 \rightarrow 18$
 $k = -16 \rightarrow 16$
 $l = -8 \rightarrow 8$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.134$
 $S = 0.94$
2154 reflections
199 parameters
0 restraints
11 constraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0688P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.19 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.22 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	1.01380 (5)	0.90521 (5)	0.19233 (12)	0.0434 (2)
N3	0.86107 (14)	0.80121 (14)	0.1963 (3)	0.0328 (5)
N2	0.97880 (15)	0.70583 (16)	0.2103 (4)	0.0444 (6)
N1	0.90331 (15)	0.64596 (17)	0.2066 (4)	0.0416 (6)
N4	0.80738 (14)	0.88156 (15)	0.1519 (3)	0.0330 (5)
C10	0.83938 (16)	0.96827 (18)	0.1890 (4)	0.0321 (6)
C7	0.83399 (17)	0.70319 (18)	0.1968 (4)	0.0340 (6)
C6	0.74169 (17)	0.66820 (19)	0.1859 (4)	0.0351 (6)
C9	0.92724 (16)	0.9844 (2)	0.2857 (4)	0.0391 (6)
H9A	0.9215	0.9715	0.4202	0.047*
H9B	0.9444	1.0533	0.2709	0.047*
C11	0.78112 (17)	1.05326 (18)	0.1437 (4)	0.0343 (6)
C8	0.95214 (16)	0.79732 (19)	0.2038 (4)	0.0357 (6)
F1	0.61197 (13)	1.28437 (13)	0.0296 (4)	0.0784 (7)
C5	0.7266 (2)	0.5726 (2)	0.1183 (4)	0.0450 (7)

H5	0.7739	0.5326	0.0841	0.054*
C15	0.6392 (2)	1.1137 (2)	0.0357 (5)	0.0543 (8)
H15	0.5822	1.1019	-0.0114	0.065*
C1	0.67026 (18)	0.7263 (2)	0.2400 (5)	0.0432 (7)
H1	0.6800	0.7897	0.2881	0.052*
C16	0.69570 (18)	1.0365 (2)	0.0751 (5)	0.0458 (7)
H16	0.6764	0.9716	0.0556	0.055*
C2	0.5845 (2)	0.6898 (2)	0.2223 (6)	0.0571 (9)
H2	0.5366	0.7289	0.2570	0.068*
C12	0.80794 (19)	1.1509 (2)	0.1700 (5)	0.0473 (7)
H12	0.8653	1.1640	0.2139	0.057*
C3	0.5708 (2)	0.5959 (2)	0.1535 (6)	0.0597 (9)
H3	0.5133	0.5714	0.1412	0.072*
C14	0.6683 (2)	1.2082 (2)	0.0669 (5)	0.0510 (8)
C13	0.7514 (2)	1.2285 (2)	0.1325 (6)	0.0559 (9)
H13	0.7697	1.2938	0.1517	0.067*
C4	0.6409 (2)	0.5377 (2)	0.1024 (5)	0.0583 (9)
H4	0.6306	0.4739	0.0566	0.070*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0270 (3)	0.0420 (4)	0.0614 (5)	-0.0019 (3)	0.0009 (3)	0.0042 (3)
N3	0.0284 (11)	0.0298 (11)	0.0401 (12)	0.0015 (9)	-0.0039 (9)	-0.0011 (9)
N2	0.0332 (12)	0.0383 (13)	0.0616 (16)	0.0045 (10)	-0.0029 (11)	0.0027 (11)
N1	0.0375 (12)	0.0335 (12)	0.0535 (14)	0.0038 (10)	-0.0051 (11)	0.0004 (11)
N4	0.0289 (11)	0.0295 (11)	0.0406 (12)	0.0041 (9)	-0.0034 (9)	-0.0001 (9)
C10	0.0302 (12)	0.0313 (13)	0.0349 (13)	-0.0028 (10)	0.0005 (11)	-0.0021 (11)
C7	0.0375 (14)	0.0303 (13)	0.0340 (13)	0.0012 (11)	-0.0024 (11)	-0.0027 (10)
C6	0.0353 (13)	0.0323 (13)	0.0376 (14)	-0.0041 (11)	-0.0029 (11)	0.0018 (11)
C9	0.0326 (13)	0.0348 (14)	0.0494 (16)	-0.0012 (11)	-0.0055 (12)	-0.0004 (12)
C11	0.0322 (13)	0.0318 (13)	0.0387 (14)	0.0002 (11)	-0.0001 (11)	-0.0007 (11)
C8	0.0286 (13)	0.0366 (14)	0.0418 (15)	0.0009 (11)	-0.0015 (11)	0.0019 (11)
F1	0.0629 (12)	0.0484 (11)	0.123 (2)	0.0259 (9)	-0.0115 (13)	0.0061 (12)
C5	0.0494 (17)	0.0335 (14)	0.0520 (18)	-0.0018 (13)	0.0006 (14)	-0.0031 (13)
C15	0.0394 (16)	0.0508 (18)	0.072 (2)	0.0069 (14)	-0.0123 (16)	-0.0005 (16)
C1	0.0391 (14)	0.0401 (15)	0.0504 (17)	-0.0038 (12)	-0.0037 (13)	-0.0075 (13)
C16	0.0376 (15)	0.0344 (14)	0.065 (2)	0.0021 (12)	-0.0100 (15)	-0.0013 (13)
C2	0.0392 (16)	0.059 (2)	0.073 (2)	-0.0003 (15)	-0.0022 (16)	-0.0074 (17)
C12	0.0388 (15)	0.0372 (15)	0.0656 (19)	0.0006 (13)	-0.0069 (14)	-0.0013 (15)
C3	0.0426 (17)	0.061 (2)	0.075 (2)	-0.0164 (15)	-0.0093 (17)	-0.0050 (18)
C14	0.0467 (16)	0.0428 (16)	0.064 (2)	0.0178 (14)	0.0001 (15)	0.0044 (15)
C13	0.0521 (17)	0.0326 (15)	0.083 (2)	0.0020 (13)	-0.0015 (18)	-0.0027 (15)
C4	0.061 (2)	0.0440 (17)	0.070 (2)	-0.0170 (16)	-0.0060 (18)	-0.0085 (16)

Geometric parameters (Å, °)

S1—C8	1.728 (3)	F1—C14	1.354 (3)
S1—C9	1.819 (3)	C5—C4	1.378 (4)
N3—C8	1.375 (3)	C5—H5	0.9300
N3—C7	1.382 (3)	C15—C14	1.362 (4)
N3—N4	1.384 (3)	C15—C16	1.370 (4)
N2—C8	1.296 (3)	C15—H15	0.9300
N2—N1	1.395 (3)	C1—C2	1.388 (4)
N1—C7	1.300 (3)	C1—H1	0.9300
N4—C10	1.288 (3)	C16—H16	0.9300
C10—C11	1.474 (3)	C2—C3	1.369 (4)
C10—C9	1.495 (3)	C2—H2	0.9300
C7—C6	1.471 (4)	C12—C13	1.371 (4)
C6—C1	1.391 (4)	C12—H12	0.9300
C6—C5	1.390 (4)	C3—C4	1.371 (5)
C9—H9A	0.9700	C3—H3	0.9300
C9—H9B	0.9700	C14—C13	1.356 (5)
C11—C16	1.387 (4)	C13—H13	0.9300
C11—C12	1.387 (4)	C4—H4	0.9300
C8—S1—C9	94.87 (13)	C4—C5—H5	120.3
C8—N3—C7	105.0 (2)	C6—C5—H5	120.3
C8—N3—N4	128.2 (2)	C14—C15—C16	118.6 (3)
C7—N3—N4	125.1 (2)	C14—C15—H15	120.7
C8—N2—N1	107.2 (2)	C16—C15—H15	120.7
C7—N1—N2	108.3 (2)	C6—C1—C2	120.2 (3)
C10—N4—N3	116.5 (2)	C6—C1—H1	119.9
N4—C10—C11	116.1 (2)	C2—C1—H1	119.9
N4—C10—C9	123.4 (2)	C15—C16—C11	121.2 (3)
C11—C10—C9	120.4 (2)	C15—C16—H16	119.4
N1—C7—N3	109.2 (2)	C11—C16—H16	119.4
N1—C7—C6	125.0 (2)	C3—C2—C1	119.6 (3)
N3—C7—C6	125.8 (2)	C3—C2—H2	120.2
C1—C6—C5	119.5 (3)	C1—C2—H2	120.2
C1—C6—C7	122.9 (2)	C13—C12—C11	121.1 (3)
C5—C6—C7	117.6 (2)	C13—C12—H12	119.4
C10—C9—S1	112.70 (19)	C11—C12—H12	119.4
C10—C9—H9A	109.1	C2—C3—C4	120.6 (3)
S1—C9—H9A	109.1	C2—C3—H3	119.7
C10—C9—H9B	109.1	C4—C3—H3	119.7
S1—C9—H9B	109.1	F1—C14—C13	119.1 (3)
H9A—C9—H9B	107.8	F1—C14—C15	118.5 (3)
C16—C11—C12	117.9 (3)	C13—C14—C15	122.4 (3)
C16—C11—C10	119.7 (2)	C14—C13—C12	118.7 (3)
C12—C11—C10	122.4 (2)	C14—C13—H13	120.6
N2—C8—N3	110.3 (2)	C12—C13—H13	120.6
N2—C8—S1	129.3 (2)	C3—C4—C5	120.7 (3)

N3—C8—S1	120.31 (19)	C3—C4—H4	119.6
C4—C5—C6	119.4 (3)	C5—C4—H4	119.6
C8—N2—N1—C7	0.4 (3)	C7—N3—C8—N2	-0.6 (3)
C8—N3—N4—C10	-28.0 (4)	N4—N3—C8—N2	-166.4 (3)
C7—N3—N4—C10	168.8 (2)	C7—N3—C8—S1	176.35 (19)
N3—N4—C10—C11	-179.2 (2)	N4—N3—C8—S1	10.5 (4)
N3—N4—C10—C9	-3.7 (4)	C9—S1—C8—N2	-158.0 (3)
N2—N1—C7—N3	-0.8 (3)	C9—S1—C8—N3	25.7 (2)
N2—N1—C7—C6	178.6 (3)	C1—C6—C5—C4	1.3 (5)
C8—N3—C7—N1	0.9 (3)	C7—C6—C5—C4	-178.8 (3)
N4—N3—C7—N1	167.3 (2)	C5—C6—C1—C2	-1.5 (5)
C8—N3—C7—C6	-178.5 (3)	C7—C6—C1—C2	178.6 (3)
N4—N3—C7—C6	-12.1 (4)	C14—C15—C16—C11	-0.6 (5)
N1—C7—C6—C1	157.2 (3)	C12—C11—C16—C15	-0.4 (5)
N3—C7—C6—C1	-23.5 (4)	C10—C11—C16—C15	178.5 (3)
N1—C7—C6—C5	-22.7 (4)	C6—C1—C2—C3	0.8 (5)
N3—C7—C6—C5	156.5 (3)	C16—C11—C12—C13	1.1 (5)
N4—C10—C9—S1	44.7 (3)	C10—C11—C12—C13	-177.9 (3)
C11—C10—C9—S1	-140.1 (2)	C1—C2—C3—C4	0.2 (6)
C8—S1—C9—C10	-48.7 (2)	C16—C15—C14—F1	-179.3 (3)
N4—C10—C11—C16	3.3 (4)	C16—C15—C14—C13	1.1 (6)
C9—C10—C11—C16	-172.2 (3)	F1—C14—C13—C12	180.0 (3)
N4—C10—C11—C12	-177.8 (3)	C15—C14—C13—C12	-0.5 (6)
C9—C10—C11—C12	6.7 (4)	C11—C12—C13—C14	-0.6 (5)
N1—N2—C8—N3	0.1 (3)	C2—C3—C4—C5	-0.4 (6)
N1—N2—C8—S1	-176.5 (2)	C6—C5—C4—C3	-0.4 (5)

Hydrogen-bond geometry (Å, °)

Cg3 is the centroid of the C1—C6 phenyl ring.

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C9—H9B...N2 ⁱ	0.97	2.36	3.301 (3)	164
C12—H12...N2 ⁱ	0.93	2.47	3.393 (4)	173
C4—H4...F1 ⁱⁱ	0.93	2.57	3.475 (4)	164
C1—H1...Cg3 ⁱⁱⁱ	0.93	2.93	3.598 (3)	130

Symmetry codes: (i) $-x+2, y+1/2, -z+1/2$; (ii) $x, y-1, z$; (iii) $x, -y+3/2, z+1/2$.