

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3,5-Bis(4-bromophenyl)-1-phenyl-4,5-dihydro-1H-pyrazole

 S. Samshuddin,^a B. Narayana,^a H. S. Yathirajan,^{b‡}
 A. P. Safwan^c and Edward R. T. Tiekink^{c*}
^aDepartment of Studies in Chemistry, Mangalore University, Mangalagangotri 574 199, India, ^bDepartment of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore 570 006, India, and ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: edward.tiekink@gmail.com


Received 28 April 2010; accepted 29 April 2010

 Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.027; wR factor = 0.052; data-to-parameter ratio = 18.6.

In the title compound, $\text{C}_{21}\text{H}_{16}\text{Br}_2\text{N}_2$, the central pyrazole ring adopts an flattened envelope conformation, with the stereogenic C atom in the flap position. The deviations from planarity for this ring are relatively minor (r.m.s. deviation = 0.045 Å) and the dihedral angles formed with the N- and C_{imine}-bound benzene rings are 7.73 (13) and 11.00 (13)°, respectively. By contrast, the benzene ring bound at the chiral C atom is almost orthogonal to the rest of the molecule; the dihedral angle formed between this ring and the pyrazole ring is 79.53 (13)°. In the crystal, the packing is stabilized by C—H...N and C—H...Br interactions.

Related literature

For the pharmacological activity of pyrazoline derivatives, see: Hes *et al.* (1978); Amir *et al.* (2008); Sarojini *et al.* (2010). For related structures, see: Fun *et al.* (2010); Yathirajan *et al.* (2007). For the structure of the parent compound, 1,3,5-triphenyl-2-pyrazoline, see: Foces-Foces *et al.* (2001). For conformational analysis, see: Cremer & Pople (1975).


[‡] Additional correspondence author, e-mail: yathirajan@hotmail.com.

Experimental

Crystal data

$\text{C}_{21}\text{H}_{16}\text{Br}_2\text{N}_2$	$V = 1833.79$ (9) Å ³
$M_r = 456.18$	$Z = 4$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation
$a = 10.5815$ (3) Å	$\mu = 4.43$ mm ⁻¹
$b = 11.2119$ (3) Å	$T = 100$ K
$c = 15.4569$ (4) Å	$0.35 \times 0.15 \times 0.08$ mm

Data collection

Bruker SMART APEX diffractometer	17504 measured reflections
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	4201 independent reflections
$T_{\min} = 0.537$, $T_{\max} = 0.746$	3753 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.044$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.027$	$\Delta\rho_{\text{max}} = 0.60$ e Å ⁻³
$wR(F^2) = 0.052$	$\Delta\rho_{\text{min}} = -0.40$ e Å ⁻³
$S = 1.02$	Absolute structure: Flack (1983),
4201 reflections	1804 Friedel pairs
226 parameters	Flack parameter: 0.003 (7)
H-atom parameters constrained	

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C5}-\text{H5}\cdots\text{N1}^i$	0.95	2.58	3.374 (3)	141
$\text{C20}-\text{H20}\cdots\text{Br1}^{ii}$	0.95	2.92	3.768 (2)	148

 Symmetry codes: (i) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x, y - 1, z$.

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SIR97 (Altomare *et al.*, 1999); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and DIAMOND (Brandenburg, 2006); software used to prepare material for publication: publCIF (Westrip, 2010).

SS thanks Mangalore University and the UGC SAP for financial assistance for the purchase of chemicals. HSY thanks the University of Mysore for the sanction of sabbatical leave

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5428).

References

- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Amir, M., Kumar, H. & Khan, S. A. (2008). *Bioorg. Med. Chem. Lett.* **18**, 918–922.
- Brandenburg, K. (2006). DIAMOND. Crystal Impact GbR, Bonn, Germany.
- Bruker (2008). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Foces-Foces, C., Jagerovic, N. & Elguero, J. (2001). *Z. Kristallogr.* **216**, 240–244.
- Fun, H.-K., Hemamalini, M., Samshuddin, S., Narayana, B. & Yathirajan, H. S. (2010). *Acta Cryst.* **E66**, o582–o583.
- Hes, R. V., Wellinga, K. & Grosscurt, A. C. (1978). *J. Agric. Food Chem.* **26**, 915–918.

- Sarojini, B. K., Vidyagayatri, M., Darshanraj, C. G., Bharath, B. R. & Manjunatha, H. (2010). *Lett. Drug Design Discov.* **7**, 214–224.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**. Submitted.
- Yathirajan, H. S., Bindya, S., Sarojini, B. K., Narayana, B. & Bolte, M. (2007). *Acta Cryst.* **E63**, o2718.

supporting information

Acta Cryst. (2010). E66, o1279–o1280 [https://doi.org/10.1107/S1600536810015795]

3,5-Bis(4-bromophenyl)-1-phenyl-4,5-dihydro-1H-pyrazole

S. Samshuddin, B. Narayana, H. S. Yathirajan, A. P. Safwan and Edward R. T. Tiekink

S1. Comment

Derivatives of pyrazoline possess a range of pharmacological activities, having, for example, anti-tumour, anti-microbial, and anti-tubercular activities (Hes *et al.*, 1978; Amir *et al.*, 2008). Further, some of these compounds also have anti-inflammatory, anti-diabetic, anaesthetic, analgesic and DPPH scavenging properties (Sarojini *et al.*, 2010). In continuation of previous structural studies of pyrazoline derivatives (Fun *et al.*, 2010, Yathirajan *et al.*, 2007), the title compound, (I), was synthesised and its crystal structure determined.

The structure analysis of (I) shows the C1 centre to have an *S* configuration, Fig. 1. The conformation of the central pyrazole ring is an envelope on the C1 atom as defined by the ring-puckering parameters of $q_2 = 0.101(3) \text{ \AA}$ and $\varphi_2 = 251.7(14)^\circ$ (Cremer & Pople, 1975). That being stated, the maximum deviations from the five atoms of the ring are 0.052(2) and -0.064(3) \AA for the N2 and C1 atoms, respectively; the r.m.s. deviation = 0.0450 \AA . The N2- and C3-bound benzene rings are approximately co-planar with the central ring as seen in the dihedral angles formed between their respective least-squares planes and that through the pyrazole ring of 7.73(13) and 11.00(13) $^\circ$; the dihedral angle between these benzene rings is 4.32(12) $^\circ$. By contrast, the C1-bound benzene ring is almost orthogonal to the remaining molecule with a dihedral angle of 79.53(13) $^\circ$ formed between it and the pyrazole ring. To a first approximation, the overall conformation in (I) resembles that in the analogous 1,3,5-triphenyl-2-pyrazoline "parent" compound although the deviations from planarity are slightly greater in the literature structure (Foces-Foces *et al.*, 2001). Further, the N1–N2 [1.369(3) \AA] and N1=C3 [1.291(3) \AA] bond distances in (I) are comparable to the equivalent distances in 1,3,5-triphenyl-2-pyrazoline of 1.387(5) and 1.285(7) \AA , respectively.

The molecules are consolidated into a 3-D network by C–H \cdots N and C–H \cdots Br contacts, Fig. 2 and Table 1.

S2. Experimental

A mixture of (2*E*)-1,3-bis(4-bromophenyl)prop-2-en-1-one (3.66 g, 0.01 mol) and phenyl hydrazine (1.08 g, 0.01 mol) in glacial acetic acid (50 ml) was refluxed for 6 h. The reaction mixture was cooled and poured into ice-cold water (50 ml). The precipitate was collected by filtration and purified by recrystallization from ethanol. Yellow blocks of (I) were grown from toluene by slow evaporation; the yield was 86%, m.pt. 481 K. Analytical data (%): Found (Calc'd): C 55.21 (55.29); H 3.48 (3.54); N 6.10 (6.14).

S3. Refinement

Carbon-bound H-atoms were placed in calculated positions (C–H 0.95 to 1.00 \AA) and were included in the refinement in the riding model approximation, with $U_{\text{iso}}(\text{H})$ set to 1.2 to 1.5 $U_{\text{eq}}(\text{C})$.




Figure 1

The molecular structure of (I) showing displacement ellipsoids at the 50% probability level.




Figure 2

A view in projection down the *a* axis of the crystal packing in (I) mediated by C–H···N and C–H···Br contacts, shown as orange and purple dashed lines, respectively.

3,5-Bis(4-bromophenyl)-1-phenyl-4,5-dihydro-1*H*-pyrazole

Crystal data

$C_{21}H_{16}Br_2N_2$

$M_r = 456.18$

Orthorhombic, $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 10.5815(3) \text{ \AA}$

$b = 11.2119(3) \text{ \AA}$

$c = 15.4569 (4) \text{ \AA}$
 $V = 1833.79 (9) \text{ \AA}^3$
 $Z = 4$
 $F(000) = 904$
 $D_x = 1.652 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 4444 reflections

$\theta = 2.3\text{--}26.2^\circ$
 $\mu = 4.43 \text{ mm}^{-1}$
 $T = 100 \text{ K}$
 Block, yellow
 $0.35 \times 0.15 \times 0.08 \text{ mm}$

Data collection

Bruker SMART APEX
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 ω scans
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.537$, $T_{\max} = 0.746$

17504 measured reflections
 4201 independent reflections
 3753 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.044$
 $\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 2.2^\circ$
 $h = -13 \rightarrow 13$
 $k = -14 \rightarrow 14$
 $l = -20 \rightarrow 20$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.027$
 $wR(F^2) = 0.052$
 $S = 1.02$
 4201 reflections
 226 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map

Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0105P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.60 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.40 \text{ e \AA}^{-3}$
 Absolute structure: Flack (1983), 1804 Friedel
 pairs
 Absolute structure parameter: 0.003 (7)

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.45694 (3)	0.39983 (2)	-0.095975 (18)	0.03237 (9)
Br2	1.28094 (3)	-0.06644 (3)	0.379191 (19)	0.02784 (8)
N1	0.68637 (18)	-0.10495 (19)	0.19763 (12)	0.0155 (5)
N2	0.56377 (19)	-0.07418 (19)	0.18013 (13)	0.0175 (5)
C1	0.5321 (3)	0.0486 (2)	0.20739 (15)	0.0184 (5)
H1	0.4556	0.0479	0.2453	0.022*
C2	0.6507 (2)	0.0809 (3)	0.26139 (16)	0.0226 (6)
H2A	0.6878	0.1576	0.2422	0.027*
H2B	0.6304	0.0859	0.3238	0.027*

C3	0.7385 (2)	-0.0217 (2)	0.24286 (16)	0.0144 (6)
C4	0.5117 (2)	0.1321 (2)	0.13185 (17)	0.0165 (6)
C5	0.4266 (3)	0.2261 (2)	0.13916 (17)	0.0206 (6)
H5	0.3790	0.2351	0.1909	0.025*
C6	0.4100 (3)	0.3064 (2)	0.07249 (17)	0.0230 (6)
H6	0.3521	0.3707	0.0781	0.028*
C7	0.4791 (3)	0.2917 (2)	-0.00225 (17)	0.0208 (6)
C8	0.5631 (3)	0.1994 (2)	-0.01219 (16)	0.0213 (6)
H8	0.6087	0.1901	-0.0647	0.026*
C9	0.5803 (2)	0.1200 (2)	0.05566 (17)	0.0193 (6)
H9	0.6394	0.0567	0.0500	0.023*
C10	0.8685 (2)	-0.0309 (2)	0.27394 (16)	0.0141 (6)
C11	0.9485 (3)	-0.1218 (2)	0.24481 (16)	0.0179 (6)
H11	0.9182	-0.1784	0.2041	0.021*
C12	1.0714 (3)	-0.1300 (2)	0.27476 (17)	0.0199 (6)
H12	1.1256	-0.1914	0.2544	0.024*
C13	1.1146 (2)	-0.0478 (3)	0.33476 (17)	0.0190 (6)
C14	1.0378 (2)	0.0425 (2)	0.36482 (16)	0.0190 (5)
H14	1.0685	0.0981	0.4062	0.023*
C15	0.9159 (2)	0.0512 (2)	0.33413 (16)	0.0178 (6)
H15	0.8631	0.1140	0.3541	0.021*
C16	0.4888 (2)	-0.1457 (2)	0.12849 (16)	0.0168 (5)
C17	0.3627 (2)	-0.1161 (2)	0.11246 (16)	0.0195 (6)
H17	0.3271	-0.0468	0.1380	0.023*
C18	0.2893 (3)	-0.1876 (3)	0.05953 (17)	0.0223 (6)
H18	0.2036	-0.1666	0.0494	0.027*
C19	0.3379 (3)	-0.2884 (3)	0.02115 (17)	0.0222 (6)
H19	0.2869	-0.3362	-0.0157	0.027*
C20	0.4627 (3)	-0.3189 (2)	0.03732 (17)	0.0225 (6)
H20	0.4969	-0.3886	0.0113	0.027*
C21	0.5387 (3)	-0.2497 (2)	0.09078 (16)	0.0188 (5)
H21	0.6236	-0.2725	0.1018	0.023*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0559 (2)	0.01873 (14)	0.02246 (14)	0.00349 (14)	-0.01182 (14)	0.00164 (12)
Br2	0.01697 (13)	0.03195 (17)	0.03461 (17)	0.00011 (12)	-0.00644 (13)	0.00337 (14)
N1	0.0142 (11)	0.0171 (11)	0.0151 (10)	-0.0003 (9)	-0.0004 (8)	0.0027 (10)
N2	0.0142 (11)	0.0189 (12)	0.0193 (10)	0.0029 (10)	-0.0035 (9)	-0.0018 (9)
C1	0.0191 (14)	0.0189 (13)	0.0172 (13)	0.0012 (12)	0.0014 (11)	-0.0035 (11)
C2	0.0227 (14)	0.0244 (16)	0.0208 (13)	0.0058 (13)	-0.0072 (12)	-0.0030 (13)
C3	0.0172 (14)	0.0140 (13)	0.0121 (12)	-0.0011 (10)	0.0006 (10)	0.0028 (10)
C4	0.0134 (13)	0.0182 (13)	0.0179 (13)	-0.0005 (10)	-0.0044 (11)	-0.0022 (11)
C5	0.0194 (14)	0.0231 (14)	0.0195 (14)	0.0009 (11)	-0.0028 (11)	-0.0035 (12)
C6	0.0255 (16)	0.0166 (14)	0.0269 (16)	0.0043 (12)	-0.0055 (13)	-0.0047 (12)
C7	0.0284 (17)	0.0171 (14)	0.0170 (13)	-0.0012 (12)	-0.0115 (12)	-0.0024 (12)
C8	0.0264 (17)	0.0217 (15)	0.0159 (13)	-0.0010 (12)	-0.0008 (12)	-0.0018 (11)

C9	0.0176 (14)	0.0183 (15)	0.0219 (13)	0.0058 (11)	-0.0012 (11)	-0.0030 (12)
C10	0.0150 (14)	0.0137 (13)	0.0136 (12)	-0.0018 (10)	0.0004 (10)	0.0022 (10)
C11	0.0209 (15)	0.0173 (14)	0.0156 (12)	-0.0048 (12)	-0.0016 (12)	-0.0017 (10)
C12	0.0203 (15)	0.0163 (14)	0.0232 (14)	0.0003 (11)	0.0030 (12)	0.0011 (11)
C13	0.0124 (13)	0.0239 (15)	0.0205 (13)	-0.0005 (11)	-0.0010 (11)	0.0094 (12)
C14	0.0222 (14)	0.0176 (13)	0.0173 (13)	-0.0044 (11)	-0.0033 (12)	0.0025 (11)
C15	0.0210 (14)	0.0143 (13)	0.0180 (13)	0.0003 (11)	0.0037 (11)	0.0002 (11)
C16	0.0189 (13)	0.0188 (12)	0.0126 (12)	-0.0049 (10)	0.0008 (11)	0.0013 (11)
C17	0.0218 (14)	0.0186 (13)	0.0181 (14)	-0.0005 (11)	-0.0008 (12)	0.0025 (12)
C18	0.0187 (15)	0.0292 (16)	0.0190 (14)	-0.0032 (13)	-0.0030 (13)	0.0058 (12)
C19	0.0255 (16)	0.0243 (16)	0.0168 (14)	-0.0098 (13)	-0.0023 (12)	0.0001 (12)
C20	0.0249 (16)	0.0230 (14)	0.0196 (14)	-0.0061 (13)	0.0040 (13)	-0.0033 (11)
C21	0.0159 (14)	0.0221 (13)	0.0183 (13)	-0.0028 (11)	0.0021 (12)	0.0021 (12)

Geometric parameters (Å, °)

Br1—C7	1.903 (3)	C9—H9	0.9500
Br2—C13	1.901 (2)	C10—C11	1.399 (4)
N1—C3	1.291 (3)	C10—C15	1.402 (4)
N1—N2	1.369 (3)	C11—C12	1.383 (4)
N2—C16	1.382 (3)	C11—H11	0.9500
N2—C1	1.478 (3)	C12—C13	1.385 (4)
C1—C4	1.512 (4)	C12—H12	0.9500
C1—C2	1.550 (4)	C13—C14	1.379 (4)
C1—H1	1.0000	C14—C15	1.377 (4)
C2—C3	1.506 (4)	C14—H14	0.9500
C2—H2A	0.9900	C15—H15	0.9500
C2—H2B	0.9900	C16—C17	1.397 (3)
C3—C10	1.461 (4)	C16—C21	1.407 (3)
C4—C9	1.390 (4)	C17—C18	1.383 (4)
C4—C5	1.391 (4)	C17—H17	0.9500
C5—C6	1.380 (4)	C18—C19	1.377 (4)
C5—H5	0.9500	C18—H18	0.9500
C6—C7	1.377 (4)	C19—C20	1.386 (4)
C6—H6	0.9500	C19—H19	0.9500
C7—C8	1.373 (4)	C20—C21	1.390 (4)
C8—C9	1.388 (4)	C20—H20	0.9500
C8—H8	0.9500	C21—H21	0.9500
C3—N1—N2	109.2 (2)	C11—C10—C15	118.4 (2)
N1—N2—C16	120.8 (2)	C11—C10—C3	121.0 (2)
N1—N2—C1	113.1 (2)	C15—C10—C3	120.6 (2)
C16—N2—C1	125.1 (2)	C12—C11—C10	120.6 (2)
N2—C1—C4	112.9 (2)	C12—C11—H11	119.7
N2—C1—C2	100.8 (2)	C10—C11—H11	119.7
C4—C1—C2	112.8 (2)	C11—C12—C13	119.3 (2)
N2—C1—H1	110.0	C11—C12—H12	120.3
C4—C1—H1	110.0	C13—C12—H12	120.3

C2—C1—H1	110.0	C14—C13—C12	121.3 (2)
C3—C2—C1	102.6 (2)	C14—C13—Br2	120.3 (2)
C3—C2—H2A	111.2	C12—C13—Br2	118.3 (2)
C1—C2—H2A	111.2	C15—C14—C13	119.2 (2)
C3—C2—H2B	111.2	C15—C14—H14	120.4
C1—C2—H2B	111.2	C13—C14—H14	120.4
H2A—C2—H2B	109.2	C14—C15—C10	121.1 (2)
N1—C3—C10	122.0 (2)	C14—C15—H15	119.4
N1—C3—C2	113.1 (2)	C10—C15—H15	119.4
C10—C3—C2	124.9 (2)	N2—C16—C17	120.9 (2)
C9—C4—C5	118.8 (2)	N2—C16—C21	120.3 (2)
C9—C4—C1	121.3 (2)	C17—C16—C21	118.8 (2)
C5—C4—C1	119.9 (2)	C18—C17—C16	120.3 (3)
C6—C5—C4	121.0 (2)	C18—C17—H17	119.9
C6—C5—H5	119.5	C16—C17—H17	119.9
C4—C5—H5	119.5	C19—C18—C17	121.4 (3)
C7—C6—C5	118.8 (2)	C19—C18—H18	119.3
C7—C6—H6	120.6	C17—C18—H18	119.3
C5—C6—H6	120.6	C18—C19—C20	118.7 (3)
C8—C7—C6	121.8 (2)	C18—C19—H19	120.6
C8—C7—Br1	118.3 (2)	C20—C19—H19	120.6
C6—C7—Br1	119.8 (2)	C19—C20—C21	121.4 (3)
C7—C8—C9	118.9 (2)	C19—C20—H20	119.3
C7—C8—H8	120.5	C21—C20—H20	119.3
C9—C8—H8	120.5	C20—C21—C16	119.5 (3)
C8—C9—C4	120.6 (2)	C20—C21—H21	120.3
C8—C9—H9	119.7	C16—C21—H21	120.3
C4—C9—H9	119.7		
C3—N1—N2—C16	176.5 (2)	N1—C3—C10—C11	10.1 (4)
C3—N1—N2—C1	7.3 (3)	C2—C3—C10—C11	-171.9 (2)
N1—N2—C1—C4	110.1 (2)	N1—C3—C10—C15	-169.7 (2)
C16—N2—C1—C4	-58.6 (3)	C2—C3—C10—C15	8.4 (4)
N1—N2—C1—C2	-10.5 (3)	C15—C10—C11—C12	-0.1 (4)
C16—N2—C1—C2	-179.1 (2)	C3—C10—C11—C12	-179.8 (2)
N2—C1—C2—C3	9.2 (2)	C10—C11—C12—C13	0.6 (4)
C4—C1—C2—C3	-111.5 (2)	C11—C12—C13—C14	-0.4 (4)
N2—N1—C3—C10	178.0 (2)	C11—C12—C13—Br2	176.76 (19)
N2—N1—C3—C2	-0.3 (3)	C12—C13—C14—C15	-0.3 (4)
C1—C2—C3—N1	-6.1 (3)	Br2—C13—C14—C15	-177.42 (19)
C1—C2—C3—C10	175.7 (2)	C13—C14—C15—C10	0.8 (4)
N2—C1—C4—C9	-34.0 (3)	C11—C10—C15—C14	-0.7 (4)
C2—C1—C4—C9	79.5 (3)	C3—C10—C15—C14	179.1 (2)
N2—C1—C4—C5	148.6 (2)	N1—N2—C16—C17	178.2 (2)
C2—C1—C4—C5	-98.0 (3)	C1—N2—C16—C17	-13.9 (4)
C9—C4—C5—C6	-0.2 (4)	N1—N2—C16—C21	-1.9 (3)
C1—C4—C5—C6	177.3 (2)	C1—N2—C16—C21	165.9 (2)
C4—C5—C6—C7	0.4 (4)	N2—C16—C17—C18	179.0 (2)

C5—C6—C7—C8	0.2 (4)	C21—C16—C17—C18	-0.8 (4)
C5—C6—C7—Br1	179.2 (2)	C16—C17—C18—C19	-0.3 (4)
C6—C7—C8—C9	-1.1 (4)	C17—C18—C19—C20	0.9 (4)
Br1—C7—C8—C9	179.9 (2)	C18—C19—C20—C21	-0.3 (4)
C7—C8—C9—C4	1.3 (4)	C19—C20—C21—C16	-0.8 (4)
C5—C4—C9—C8	-0.6 (4)	N2—C16—C21—C20	-178.4 (2)
C1—C4—C9—C8	-178.1 (2)	C17—C16—C21—C20	1.4 (4)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C5—H5...N1 ⁱ	0.95	2.58	3.374 (3)	141
C20—H20...Br1 ⁱⁱ	0.95	2.92	3.768 (2)	148

Symmetry codes: (i) $-x+1, y+1/2, -z+1/2$; (ii) $x, y-1, z$.