

3-(2-Chloroethyl)-2-methyl-4H-pyrido[1,2-a]pyrimidin-4-one

Jerry P. Jasinski,^a Ray J. Butcher,^{b*} Q. N. M. Hakim Al-Arique,^c H. S. Yathirajan^c and B. Narayana^d

^aDepartment of Chemistry, Keene State College, 229 Main Street, Keene, NH 03435-2001, USA, ^bDepartment of Chemistry, Howard University, 525 College Street NW, Washington DC 20059, USA, ^cDepartment of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore 570 006, India, and ^dDepartment of Studies in Chemistry, Mangalore University, Mangalagangotri 574 199, India
Correspondence e-mail: rbutcher99@yahoo.com


Received 11 June 2009; accepted 13 July 2009

Key indicators: single-crystal X-ray study; $T = 110$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.065; wR factor = 0.181; data-to-parameter ratio = 22.4.

In the title molecule, $\text{C}_{11}\text{H}_{11}\text{ClN}_2\text{O}$, the pyrido[1,2-*a*]pyrimidine ring system is planar (maximum deviation = 0.0148 Å) and the methyl C and carbonyl O atoms are nearly coplanar to it. The chloroethyl side chain is in a synclinal conformation, nearly orthogonal to the pyrimidine ring, with a dihedral angle between the chloroethyl side chain and the pyrimidine ring of 88.5 (1)°. Weak intermolecular C—H···N and C—H···Cl hydrogen bonds along with π – π interactions between the pyrimidine and pyridine rings [centroid–centroid distance is 3.538 (2) Å] form a three-dimensional network. The crystal is a racemic twin with a 0.68 (12):0.32 (12) domain ratio. MOPAC AM1 and density functional theory (DFT) theoretical calculations at the B3-LYP/6–311+G(d,p) level support these observations.

Related literature

For related structures, see: Blaton *et al.* (1995); Chen & He (2006); Elotmani *et al.* (2002); Jottier *et al.* (1992); Koval'chukova *et al.* (2004); Peeters *et al.* (1993); Ravikumar & Sridhar, (2006); Yu *et al.* (2007). For general background to heterofused pyrimidines, see: Baraldi *et al.* (2002); Bookser *et al.* (2005); Chen *et al.* (2004); La Motta *et al.* (2007); Gabbert & Giannini (1997); Goodacre *et al.* (2006); Hossain *et al.* (1997); Joseph & Burke (1993); Nikitin & Smirnov (1994); Sabnis & Rangnekar (1990); Wang *et al.* (2004); White *et al.* (2004). For the synthesis, see: Toche *et al.* (2008). For GAUSSIAN03 theoretical calculations, see: Becke (1988, 1993); Frisch *et al.* (2004); Hehre *et al.* (1986); Lee *et al.* (1988); Schmidt & Polik (2007).



Experimental

Crystal data

$\text{C}_{11}\text{H}_{11}\text{ClN}_2\text{O}$
 $M_r = 222.67$
 Orthorhombic, $P2_12_12_1$
 $a = 4.2546$ (4) Å
 $b = 11.6274$ (10) Å
 $c = 20.604$ (2) Å
 $V = 1019.27$ (17) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.35$ mm⁻¹
 $T = 110$ K
 $0.51 \times 0.35 \times 0.12$ mm

Data collection

Oxford Diffraction Gemini R CCD diffractometer
 Absorption correction: multi-scan (CrysAlis RED; Oxford Diffraction, 2007)
 $T_{\min} = 0.835$, $T_{\max} = 0.959$
 4613 measured reflections
 3089 independent reflections
 2607 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.054$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.065$
 $wR(F^2) = 0.181$
 $S = 1.11$
 3089 reflections
 138 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.99$ e Å⁻³
 $\Delta\rho_{\min} = -0.52$ e Å⁻³
 Absolute structure: Flack (1983),
 1103 Friedel pairs
 Flack parameter: 0.32 (12)

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C5}-\text{H5A}\cdots\text{N2}^i$	0.95	2.50	3.394 (3)	157
$\text{C2}-\text{H2A}\cdots\text{Cl}^{ii}$	0.95	2.90	3.559 (3)	128

Symmetry codes: (i) $x - \frac{1}{2}, -y + \frac{3}{2}, -z + 1$; (ii) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: CrysAlisPro (Oxford Diffraction, 2007); cell refinement: CrysAlisPro; data reduction: CrysAlisPro; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

QNMHA thanks the University of Mysore for use of its research facilities. RJB acknowledges the NSF MRI program (grant No. CHE-0619278) for funds to purchase an X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2827).

References

- Baraldi, P. G., Cacciari, B., Romagnoli, R., Spalluto, G., Monopoli, A., Ongini, E., Varani, K. & Borea, P. A. (2002). *J. Med. Chem.* **45**, 115–126.
- Becke, A. D. (1988). *Phys. Rev.* **A38**, 3098–100.
- Becke, A. D. (1993). *J. Chem. Phys.* **98**, 5648–5652.
- Blaton, N. M., Peeters, O. M. & De Ranter, C. J. (1995). *Acta Cryst.* **C51**, 533–535.
- Bookser, B. C., Ugarkar, B. G., Matelich, M. C., Lemus, R. H., Alla, M., Tsuchiya, M., Nakane, M., Nagahisa, A., Wiesner, J. B. & Erion, M. D. (2005). *J. Med. Chem.* **48**, 7808–7820.
- Chen, C., Chen, C., Wilcoxon, K. M., Huang, C. Q., Xie, Y.-F., McCarthy, J. R., Webb, T. R., Zhu, Y.-F., Saunders, J., Liu, X.-J., Chen, T.-K., Bozigian, H. & Grigoriadis, D. E. (2004). *J. Med. Chem.* **47**, 4787–4798.
- Chen, H.-L. & He, H.-W. (2006). *Acta Cryst.* **E62**, o1226–o1227.
- Elotmani, B., Elmahi, M., Essassi, E. M. & Pierrot, M. (2002). *Acta Cryst.* **E58**, o388–o389.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Frisch, M. J., *et al.* (2004). *GAUSSIAN03*. Gaussian Inc., Wallingford, CT 06492, USA.
- Gabbert, J. F. & Giannini, A. J. (1997). *Am. J. Ther.* **4**, 159–164.
- Goodacre, S. C., Street, L. J., Hallett, D., Crawforth, J. M., Kelly, S., Owens, A. P., Blackaby, W. P., Lewis, R. T., Stanley, J., Smith, A. J., Ferris, P., Sohal, B., Cook, S. M., Pike, A., Brown, N., Wafford, K. A., Marshall, G., Castro, J. L. & Atack, J. R. (2006). *J. Med. Chem.* **49**, 35–38.
- Hehre, W. J., Random, L., von Schleyer, P. R. & Pople, J. A. (1986). In *Ab Initio Molecular Orbital Theory*. New York: Wiley.
- Hossain, N., Rozenski, J., De Clercq, E. & Herdewijn, P. (1997). *J. Org. Chem.* **62**, 2442–2447.
- Joseph, S. & Burke, J. M. (1993). *J. Biol. Chem.* **268**, 24515–24518.
- Jottier, W. I., De Winter, H. L., Peeters, O. M., Blaton, N. M. & De Ranter, C. J. (1992). *Acta Cryst.* **C48**, 1827–1830.
- Koval'chukova, O. V., Mordovina, N. I., Kuz'mina, N. E., Nikitin, S. V., Zaitsev, B. E., Strashnova, S. B. & Palkina, K. K. (2004). *Crystallogr. Rep.* **49**, 792–797.
- La Motta, C., Sartinit, S., Mugnaini, L., Simorini, F., Taliani, S., Salerno, S., Marini, A. M., Da Settimo, F., Lavecchia, A., Novellino, E., Cantore, M., Failli, P. & Ciuffi, M. (2007). *J. Med. Chem.* **50**, 4917–4927.
- Lee, C., Yang, W. & Parr, R. G. (1988). *Phys. Rev. B*, **37**, 785–789.
- Nikitin, S. V. & Smirnov, L. D. (1994). *Chem. Heter. Copm.* **30**, 507–522.
- Oxford Diffraction (2007). *CrysAlisPro* and *CrysAlis RED*. Oxford Diffraction Ltd, Abingdon, England.
- Peeters, O. M., Blaton, N. M. & De Ranter, C. J. (1993). *Acta Cryst.* **C49**, 1698–1700.
- Ravikumar, K. & Sridhar, B. (2006). *Acta Cryst.* **E62**, o3730–o3731.
- Sabnis, R. W. & Rangnekar, D. W. (1990). *Indian J. Technol.* **28**, 54–58.
- Schmidt, J. R. & Polik, W. F. (2007). *WebMO Pro*. WebMO, LLC: Holland, MI, USA. URL: <http://www.webmo.net>.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Toche, R. B., Ghotekar, B. K., Kazi, M. A., Patil, S. P. & Jachak, M. N. (2008). *Schol. Res. Exch.* doi:10.3814/2008/434329, 1–5.
- Wang, S., Folkes, A., Chuckowree, I., Cockcroft, X., Sohal, S., Miller, W., Milton, J., Wren, S. P., Vicker, N., Depledge, P., Scott, J., Smith, L., Jones, H., Mistry, P., Faint, R., Thompson, D. & Cocks, S. (2004). *J. Med. Chem.* **47**, 1329–1338.
- White, D. C., Greenwood, D. C., Downey, A. L., Bloomquist, J. R. & Wolfe, J. F. (2004). *Bioorg. Med. Chem.* **12**, 5711–5717.
- Yu, C.-Y., Yuan, X.-N. & Huang, Z.-T. (2007). *Acta Cryst.* **E63**, o3186.

supporting information

Acta Cryst. (2009). E65, o1987–o1988 [doi:10.1107/S1600536809027548]

3-(2-Chloroethyl)-2-methyl-4*H*-pyrido[1,2-*a*]pyrimidin-4-one

Jerry P. Jasinski, Ray J. Butcher, Q. N. M. Hakim Al-Arique, H. S. Yathirajan and B. Narayana

S1. Comment

Heterofused pyrimidines exhibit promising antiviral (Hossain *et al.* 1997), antibacterial (Sabnis & Rangnekar, 1990), anti-AIDS (Joseph & Burke, 1993), and antinociceptive (Bookser *et al.* 2005) activities. Fused pyrimidines are extensively used in neurology, particularly in the treatment of neurodegenerative disorders such as Parkinson's disease (Baraldi *et al.* 2002), antianxiety disorders (Goodacre *et al.* 2006) and depression (Chen *et al.* 2004). Fused pyrimidines are selective inhibitors for multidrug resistance (MDR) (Wang *et al.* 2004). A review on the synthesis, chemical and biological properties of pyrido[1,2-*a*]pyrimidines is described (Nikitin & Smirnov, 1994). Pyrido[1,2-*a*]pyrimidin-4-one derivatives as a novel class of selective aldose reductase inhibitors exhibiting antioxidant activity has been reported (La Motta *et al.* 2007). The synthesis and anticonvulsant evaluation of some new 2-substituted-3-arylpyrido[2,3-*d*]pyrimidinones have also been reported (White *et al.* 2004). The crystal structures of 3-{2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidino]ethyl}-6,7,8,9-tetrahydro-2-methyl-4*H*-pyrido[1,2-*a*]pyrimidin-4-one (risperidone) (Peeters *et al.* 1993), 3-{2-[4-(4-fluorobenzoyl)piperidino]ethyl}-2-methyl-4*H*-pyrido[1,2-*a*]pyrimidin-4-one (Pirenperone) (Blaton *et al.* 1995), 5-methyl-2-morpholino-3-*p*-tolyl-8,9,10,11-tetrahydro-2-benzothieno[2',3':6,5]pyrido[4,3-*d*]pyrimidin-4(3*H*)-one (Chen & He, 2006), 3-{2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)-1-piperidinyl]ethyl}-2,9-dimethyl-4*H*-pyrido[1,2-*a*]pyrimidin-4-one (Ocaperidone) (Jottier *et al.* 1992), 2-methyl-3-(3-methyl-1*H*-pyrazol-5-yl)pyrido[1,2-*a*]pyrimidin-4-one (Elotmani *et al.* 2002), 2-methyl-3-chloro-9-hydroxypyrido[1,2-*a*]pyrimidin-4-one and bis(2-methyl-3-chloro-9-hydroxypyrido[1,2-*a*]pyrimidin-4-onium) perchlorate (Koval'chukova *et al.* 2004), 3-(2-chloroethyl)-2-methyl-4-oxo-6,7,8,9-tetrahydro-4*H*-pyrido[1,2-*a*]pyrimidin-1-ium chloride (Ravikumar & Sridhar, 2006) and 9-(4-methoxybenzoyl)-1,2,3,4-tetrahydro-6*H*-pyrido[1,2-*a*]pyrimidin-6-one (Yu *et al.* 2007) have also been reported.

The title compound, (I), is an intermediate in the synthesis of risperidone, which is a potent antipsychotic agent, especially useful for treating schizophrenia (Gabbert & Giannini, 1997). In view of the importance of (I), the present paper describes its crystal structure.

The overall molecular geometry of (I), including bond distances and angles, is in good agreement with related structures (Blaton *et al.* 1995; Jottier *et al.* 1992; Peeters *et al.* 1993; Ravikumar & Sridhar, 2006). It consists of a pyridine ring fused to a substituted pyrimidine ring creating a planar ring system (maximum deviation, C1, = -0.0148 Å) with the methyl C and carbonyl O atoms nearly coplanar to the pyrimidine ring (Torsion angles C1-C9-C7-C8 = 177.6 (3)°; C2-N1-C1-O = -2.5 (4)° (Fig. 1). The sum of the angles around N1 is 360.0 (5)° indicating *sp*² hybridization. The chloroethyl side chain is in a synclinal (*-sc*) conformation (C1—C9—C10—C11 torsion angle = -86.6 (3)°), nearly orthogonal to the pyrimidine ring, with a dihedral angle separation between the C10/C11/Cl group and the pyrimidine ring of 88.5 (1)°.

While no classic hydrogen bonds are observed, a weak intermolecular hydrogen bond interaction exists between atom C5 from the pyridine ring and N2 from a nearby pyrimidine ring (Table 1 and Fig. 2). In addition, a weak intermolecular interaction between atom C2 from the pyrimidine ring and Cl from the substituted pyrimidine group also occurs, each

influencing crystal packing and, therefore, resulting in a three-dimensional network (Fig. 2). In addition, π - π interactions between N1/C1/C9/C7/N2/C6 (centroid Cg1) and N1/C2-C6 (centroid Cg2) rings of molecules at (x, y, z) and (1+x, y, z), with a Cg1...Cg2 distance of 3.538 (2) Å, provide additional stability to the crystal packing. The crystal is a racemic twin with domains of 0.68 (12) and 0.32 (12).

In support of these observations, a MOPAC AM1 (Schmidt & Polik, 2007) and density functional theory (DFT) geometry optimized theoretical calculation (Schmidt & Polik, 2007) with the *GAUSSIAN03* program package (Frisch *et al.* 2004) employing the B3-LYP (Becke 3 parameter Lee-Yang-Parr) exchange correlation functional, which combines the hybrid exchange functional of Becke (Becke, 1988, 1993) with the gradient-correlation functional of Lee, Yang and Parr (Lee *et al.* 1988) and the 6-311+G(d,p) basis set (Hehre *et al.* 1986), was performed on (I) utilizing starting geometries taken from the X-ray refinement data. In both calculations the resulting bond distances and angles remained relatively constant. However, the C9—C10—C11—Cl torsion angle decreased by 3.2 (1)° to 175.4 (3)° (MOPAC) and 0.07° to 178.5 (7)° (DFT) and the dihedral angle between the C10/C11/Cl group and the pyrimidine ring decreased by 2.3 (8)° to 86.1 (3)° (MOPAC) and by 8.3 (6)° to 80.1 (5)° (DFT), respectively.

In summary, it is clear that the collection of weak intermolecular hydrogen bond interactions and π - π intermolecular interactions do play a role in stabilizing crystal packing of (I).

S2. Experimental

The title compound was synthesized following the reported procedure (Toche *et al.* 2008). Pale yellow crystals of compound (I) were obtained by slow evaporation from ethyl acetate solution (m.p. 405–408 K). Analytical data: Found (calculated): C %: 59.28 (59.33); H%: 4.97 (4.98); N%: 12.54 (12.58).

S3. Refinement

All of the H atoms were placed in their calculated positions and then refined using the riding model with C—H = 0.95–0.99 Å, and with $U_{\text{iso}}(\text{H}) = 1.18\text{--}1.50U_{\text{eq}}(\text{C})$.




Figure 1

Molecular structure of (I), showing the atom labeling scheme and 50% probability displacement ellipsoids.




Figure 2

Packing diagram of (I), viewed down the *b* axis. Dashed lines indicate weak C5—H5A...N2 and C2—H2A...Cl intermolecular interactions.

3-(2-Chloroethyl)-2-methyl-4H-pyrido[1,2-a]pyrimidin-4-one

Crystal data

$C_{11}H_{11}ClN_2O$

$M_r = 222.67$

Orthorhombic, $P2_12_12_1$

Hall symbol: $P\ 2ac\ 2ab$

$a = 4.2546\ (4)\ \text{\AA}$

$b = 11.6274\ (10)\ \text{\AA}$

$c = 20.604\ (2)\ \text{\AA}$

$V = 1019.27\ (17)\ \text{\AA}^3$

$Z = 4$

$F(000) = 464$

$D_x = 1.451\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 2755 reflections

$\theta = 4.8\text{--}32.6^\circ$

$\mu = 0.35\ \text{mm}^{-1}$

$T = 110\ \text{K}$

Plate, colorless

$0.51 \times 0.35 \times 0.12\ \text{mm}$

Data collection

Oxford Diffraction Gemini R CCD
diffractometer

Radiation source: Enhance (Mo) X-ray Source

Graphite monochromator

Detector resolution: $10.5081\ \text{pixels mm}^{-1}$

φ and ω scans

Absorption correction: multi-scan

(*CrysAlis RED*; Oxford Diffraction, 2007)

$T_{\min} = 0.835$, $T_{\max} = 0.959$

4613 measured reflections

3089 independent reflections

2607 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.054$

$\theta_{\max} = 32.6^\circ$, $\theta_{\min} = 4.9^\circ$
 $h = -3 \rightarrow 6$

$k = -17 \rightarrow 16$
 $l = -30 \rightarrow 27$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.065$
 $wR(F^2) = 0.181$
 $S = 1.11$
 3089 reflections
 138 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map

Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.1108P)^2 + 0.2652P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.99 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.52 \text{ e } \text{\AA}^{-3}$
 Absolute structure: Flack (1983), 1103 Friedel
 pairs
 Absolute structure parameter: 0.32 (12)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl	0.49448 (19)	0.87358 (6)	0.12093 (3)	0.02056 (18)
O	0.1937 (6)	1.09071 (18)	0.30555 (10)	0.0240 (5)
N1	-0.0403 (6)	0.9961 (2)	0.39118 (10)	0.0150 (4)
N2	0.0097 (7)	0.79306 (19)	0.40322 (11)	0.0172 (4)
C1	0.1541 (7)	0.9982 (2)	0.33357 (13)	0.0151 (5)
C2	-0.1616 (9)	1.0994 (2)	0.41318 (14)	0.0207 (6)
H2A	-0.1116	1.1685	0.3909	0.025*
C3	-0.3506 (8)	1.1032 (3)	0.46596 (15)	0.0231 (6)
H3A	-0.4354	1.1744	0.4803	0.028*
C4	-0.4212 (7)	0.9999 (3)	0.49970 (14)	0.0211 (6)
H4A	-0.5530	1.0019	0.5369	0.025*
C5	-0.3006 (8)	0.8986 (2)	0.47886 (13)	0.0191 (5)
H5A	-0.3477	0.8299	0.5019	0.023*
C6	-0.1033 (7)	0.8935 (2)	0.42262 (13)	0.0147 (5)
C7	0.1973 (8)	0.7917 (2)	0.34977 (13)	0.0155 (5)
C8	0.3087 (9)	0.6735 (2)	0.33069 (16)	0.0239 (6)
H8A	0.2581	0.6188	0.3653	0.036*
H8B	0.5366	0.6750	0.3238	0.036*
H8C	0.2038	0.6499	0.2905	0.036*
C9	0.2748 (7)	0.8891 (2)	0.31499 (12)	0.0146 (5)
C10	0.4766 (9)	0.8869 (2)	0.25437 (12)	0.0182 (5)

H10A	0.6268	0.8220	0.2567	0.022*
H10B	0.5983	0.9592	0.2511	0.022*
C11	0.2671 (7)	0.8733 (3)	0.19491 (13)	0.0201 (5)
H11A	0.1130	0.9371	0.1936	0.024*
H11B	0.1489	0.8002	0.1982	0.024*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl	0.0233 (3)	0.0253 (3)	0.0132 (3)	−0.0031 (3)	0.0030 (3)	−0.0001 (2)
O	0.0323 (13)	0.0185 (9)	0.0214 (10)	−0.0029 (9)	0.0024 (10)	0.0048 (8)
N1	0.0188 (11)	0.0144 (8)	0.0118 (9)	0.0013 (9)	−0.0009 (9)	0.0005 (7)
N2	0.0200 (10)	0.0157 (8)	0.0160 (10)	−0.0002 (12)	−0.0001 (11)	0.0019 (7)
C1	0.0168 (12)	0.0163 (11)	0.0122 (11)	−0.0028 (11)	−0.0005 (10)	0.0002 (9)
C2	0.0286 (15)	0.0163 (11)	0.0173 (13)	0.0039 (12)	0.0000 (12)	−0.0001 (9)
C3	0.0259 (15)	0.0235 (13)	0.0198 (14)	0.0058 (13)	−0.0013 (12)	−0.0054 (10)
C4	0.0198 (14)	0.0300 (14)	0.0135 (12)	0.0026 (12)	0.0017 (10)	−0.0019 (10)
C5	0.0196 (13)	0.0234 (12)	0.0142 (12)	−0.0029 (12)	0.0018 (11)	0.0028 (10)
C6	0.0178 (11)	0.0146 (11)	0.0118 (11)	−0.0013 (9)	−0.0029 (9)	0.0017 (8)
C7	0.0194 (13)	0.0125 (10)	0.0146 (11)	0.0001 (11)	−0.0022 (11)	−0.0001 (9)
C8	0.0282 (16)	0.0167 (11)	0.0267 (15)	0.0015 (13)	0.0028 (14)	−0.0022 (10)
C9	0.0133 (11)	0.0183 (11)	0.0124 (12)	−0.0020 (10)	−0.0004 (9)	0.0005 (9)
C10	0.0164 (12)	0.0245 (12)	0.0138 (11)	−0.0008 (13)	0.0011 (10)	0.0003 (9)
C11	0.0178 (12)	0.0305 (13)	0.0119 (11)	−0.0035 (12)	0.0019 (10)	−0.0012 (11)

Geometric parameters (Å, °)

Cl—C11	1.805 (3)	C5—C6	1.432 (4)
O—C1	1.232 (3)	C5—H5A	0.95
N1—C2	1.383 (4)	C7—C9	1.381 (4)
N1—C6	1.384 (3)	C7—C8	1.505 (4)
N1—C1	1.447 (3)	C8—H8A	0.98
N2—C6	1.325 (3)	C8—H8B	0.98
N2—C7	1.360 (4)	C8—H8C	0.98
C1—C9	1.421 (4)	C9—C10	1.516 (4)
C2—C3	1.353 (5)	C10—C11	1.523 (4)
C2—H2A	0.95	C10—H10A	0.99
C3—C4	1.420 (5)	C10—H10B	0.99
C3—H3A	0.95	C11—H11A	0.99
C4—C5	1.355 (4)	C11—H11B	0.99
C4—H4A	0.95		
C2—N1—C6	121.5 (2)	N2—C7—C8	114.0 (2)
C2—N1—C1	117.9 (2)	C9—C7—C8	122.6 (3)
C6—N1—C1	120.6 (2)	C7—C8—H8A	109.5
C6—N2—C7	117.9 (2)	C7—C8—H8B	109.5
O—C1—C9	127.1 (3)	H8A—C8—H8B	109.5
O—C1—N1	118.5 (3)	C7—C8—H8C	109.5

C9—C1—N1	114.4 (2)	H8A—C8—H8C	109.5
C3—C2—N1	120.9 (3)	H8B—C8—H8C	109.5
C3—C2—H2A	119.5	C7—C9—C1	120.4 (2)
N1—C2—H2A	119.5	C7—C9—C10	123.3 (2)
C2—C3—C4	119.4 (3)	C1—C9—C10	116.3 (2)
C2—C3—H3A	120.3	C9—C10—C11	109.5 (3)
C4—C3—H3A	120.3	C9—C10—H10A	109.8
C5—C4—C3	120.0 (3)	C11—C10—H10A	109.8
C5—C4—H4A	120.0	C9—C10—H10B	109.8
C3—C4—H4A	120.0	C11—C10—H10B	109.8
C4—C5—C6	120.9 (3)	H10A—C10—H10B	108.2
C4—C5—H5A	119.5	C10—C11—C1	111.4 (2)
C6—C5—H5A	119.5	C10—C11—H11A	109.3
N2—C6—N1	123.3 (2)	C1—C11—H11A	109.3
N2—C6—C5	119.6 (2)	C10—C11—H11B	109.3
N1—C6—C5	117.2 (2)	C1—C11—H11B	109.3
N2—C7—C9	123.4 (2)	H11A—C11—H11B	108.0
C2—N1—C1—O	-2.5 (4)	C4—C5—C6—N2	-179.6 (3)
C6—N1—C1—O	177.1 (3)	C4—C5—C6—N1	0.5 (4)
C2—N1—C1—C9	178.7 (3)	C6—N2—C7—C9	-0.1 (4)
C6—N1—C1—C9	-1.8 (4)	C6—N2—C7—C8	-178.8 (3)
C6—N1—C2—C3	-0.9 (5)	N2—C7—C9—C1	-1.0 (4)
C1—N1—C2—C3	178.6 (3)	C8—C7—C9—C1	177.6 (3)
N1—C2—C3—C4	1.0 (5)	N2—C7—C9—C10	-178.2 (3)
C2—C3—C4—C5	-0.3 (5)	C8—C7—C9—C10	0.4 (5)
C3—C4—C5—C6	-0.5 (5)	O—C1—C9—C7	-176.9 (3)
C7—N2—C6—N1	0.2 (4)	N1—C1—C9—C7	1.8 (4)
C7—N2—C6—C5	-179.7 (3)	O—C1—C9—C10	0.6 (4)
C2—N1—C6—N2	-179.7 (3)	N1—C1—C9—C10	179.3 (2)
C1—N1—C6—N2	0.8 (4)	C7—C9—C10—C11	90.7 (3)
C2—N1—C6—C5	0.2 (4)	C1—C9—C10—C11	-86.6 (3)
C1—N1—C6—C5	-179.3 (3)	C9—C10—C11—C1	178.6 (2)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C5—H5A...N2 ⁱ	0.95	2.50	3.394 (3)	157
C2—H2A...C1 ⁱⁱ	0.95	2.90	3.559 (3)	128

Symmetry codes: (i) $x-1/2, -y+3/2, -z+1$; (ii) $-x, y+1/2, -z+1/2$.