

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# 4-[(1*E*)-3-(2,6-Dichloro-3-fluorophenyl)-3-oxoprop-1-en-1-yl]benzonitrile

 Aletti S. Praveen,<sup>a</sup> Hemmige S. Yathirajan,<sup>a</sup> Badiadka Narayana,<sup>b</sup> Thomas Gerber,<sup>c</sup> Eric Hosten<sup>c</sup> and Richard Betz<sup>c\*</sup>

<sup>a</sup>University of Mysore, Department of Studies in Chemistry, Manasagangotri, Mysore 570 006, India, <sup>b</sup>Mangalore University, Department of Studies in Chemistry, Mangalagangotri 574 199, India, and <sup>c</sup>Nelson Mandela Metropolitan University, Summerstrand Campus, Department of Chemistry, University Way, Summerstrand, PO Box 77000, Port Elizabeth 6031, South Africa  
Correspondence e-mail: richard.betz@webmail.co.za


Received 4 April 2012; accepted 10 April 2012

Key indicators: single-crystal X-ray study;  $T = 200$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å; disorder in main residue;  $R$  factor = 0.037;  $wR$  factor = 0.098; data-to-parameter ratio = 17.4.

In the title molecule,  $\text{C}_{16}\text{H}_8\text{Cl}_2\text{FNO}$ , the benzene rings form a dihedral angle of  $78.69$  (8)°. The F atom is disordered over two positions in a 0.530 (3):0.470 (3) ratio. The crystal packing exhibits  $\pi$ - $\pi$  interactions between dichloro-substituted rings [centroid-centroid distance =  $3.6671$  (10) Å] and weak intermolecular C—H...F contacts.

## Related literature

For the biological activity of chalcones, see: Rajendra Prasad *et al.* (2008); Shivakumar *et al.* (2005); Churkin *et al.* (1982); Herencia *et al.* (1998). For a related structure, see: Betz *et al.* (2012). For the graph-set analysis of hydrogen bonds, see: Etter *et al.* (1990); Bernstein *et al.* (1995).



## Experimental

## Crystal data

$\text{C}_{16}\text{H}_8\text{Cl}_2\text{FNO}$   
 $M_r = 320.13$   
 Monoclinic,  $P2_1/c$

$a = 13.2751$  (3) Å  
 $b = 8.5002$  (2) Å  
 $c = 13.9854$  (3) Å

$\beta = 116.773$  (1)°  
 $V = 1408.95$  (6) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation

$\mu = 0.47$  mm<sup>-1</sup>  
 $T = 200$  K  
 $0.55 \times 0.32 \times 0.20$  mm

## Data collection

Bruker APEXII CCD diffractometer  
 Absorption correction: multi-scan (SADABS; Bruker, 2008)  
 $T_{\min} = 0.783$ ,  $T_{\max} = 0.913$

12885 measured reflections  
 3480 independent reflections  
 2904 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.013$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$   
 $wR(F^2) = 0.098$   
 $S = 1.05$   
 3480 reflections

200 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.41$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.43$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C1}-\text{H1}\cdots\text{F1A}^i$	0.95	2.50	3.130 (2)	124

 Symmetry code: (i)  $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$ .

Data collection: APEX2 (Bruker, 2010); cell refinement: SAINT (Bruker, 2010); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and Mercury (Macrae *et al.*, 2008); software used to prepare material for publication: SHELXL97 and PLATON (Spek, 2009).

ASP thanks the University of Mysore for the research facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV5282).

## References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.  
 Betz, R., Gerber, T., Hosten, E., Praveen, A. S., Yathirajan, H. S. & Narayana, B. (2012). *Acta Cryst.* **E68**, o512.  
 Bruker (2008). *SADABS*. Bruker Inc., Madison, Wisconsin, USA.  
 Bruker (2010). *APEX2* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Churkin, Yu. D., Panfilova, L. V., Boreko, E. I., Timofeeva, M. M. & Votyakov, V. I. (1982). *Pharm. Chem. J.* **16**, 103–105.  
 Etter, M. C., MacDonald, J. C. & Bernstein, J. (1990). *Acta Cryst.* **B46**, 256–262.  
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.  
 Herencia, F., Ferrandiz, M. L., Ubada, A., Dominguez, J., Charris, J. E., Lobo, G. M. & Alcaraz, M. J. (1998). *Bioorg. Med. Chem. Lett.* **8**, 1169–1174.  
 Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.  
 Rajendra Prasad, Y., Praveen Kumar, P., Ravi Kumar, P. & Srinivas Rao, A. (2008). *E-J. Chem.* **5**, 144–148.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Shivakumar, P. M., Geetha Babu, S. M. & Mukesh, D. (2005). *Chem. Pharm. Bull.* **55**, 44–49.  
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

## supporting information

*Acta Cryst.* (2012). E68, o1413 [doi:10.1107/S1600536812015589]

**4-[(1*E*)-3-(2,6-Dichloro-3-fluorophenyl)-3-oxoprop-1-en-1-yl]benzotrile**

**Aletti S. Praveen, Hemmige S. Yathirajan, Badiadka Narayana, Thomas Gerber, Eric Hosten and Richard Betz**

**S1. Comment**

Chalcones are the products of condensation reactions of aromatic aldehydes with acetophenones in the presence of alkali. Chalcones constitute an important group of natural products, and some of them possess a wide range of biological activities such as antimicrobial (Rajendra Prasad *et al.*, 2008), antitubercular (Shivakumar *et al.*, 2005), antiviral (Churkin *et al.*, 1982) and anti-inflammatory activity (Herencia *et al.*, 1998). The crystal structures of some chalcones such as have been reported in the literature. As a part of our ongoing studies on chalcones, the title compound (I) has been synthesized and characterized by X-ray diffraction.

In (I) (Fig. 1), all bond lengths and angles are normal and correspond to those observed in (2*E*)-1-(2,6-dichloro-3-fluorophenyl)-3-(4-fluorophenyl)prop-2-en-1-one (Betz *et al.*, 2012). The fluorine atom is disordered over two positions with site occupancy factors of 0.530 (3) and 0.470 (3), respectively. The mean planes of the two aromatic rings for a dihedral angle of 78.69 (8)°.


In the crystal, C–H···F contacts (Table 1) whose range falls by more than 0.1 Å below the sum of van-der-Waals radii of the respective atoms are present. These are supported by one of the vinylic hydrogen atoms as well as the disordered fluorine atom. In terms of graph-set analysis (Etter *et al.*, 1990; Bernstein *et al.*, 1995), the descriptor for these hydrogen bonds is  $C^1_1(8)$  on the unary level. The shortest intercentroid distance between two aromatic systems was found at 3.6674 (10) Å and is apparent between the halogenated phenyl rings. The packing of the title compound in the crystal structure is shown in Figure 2.

**S2. Experimental**

To a stirred solution of 1-(2,6-dichloro-3-fluorophenyl)ethanone (1 g, 4.8 mmol) and 4-formylbenzotrile (0.62 g, 4.8 mmol) in ethanol (10 ml), powdered KOH (0.40 g, 7.2 mmol) was added at 273 K. The reaction mixture was stirred at room temperature for 2 h. After completion of the reaction, the mixture was poured into ice cold water and subsequently acidified with 1.5 N HCl (pH ~3). The precipitated solid was filtered and dried to afford 1 g of the title compound as pale yellow solid in 91% yield. The single-crystal was grown from a mixture of toluene:acetone (*v:v* = 1:1) by slow evaporation at room temperature (m.p.: 414–417 K).


**S3. Refinement**

C-bound H atoms were placed in calculated positions (C—H 0.95 Å) and were included in the refinement in the riding model approximation, with  $U(H)$  set to  $1.2U_{eq}(C)$ .



**Figure 1**

The molecular structure of the title compound showing the atomic numbering and 50% probability displacement ellipsoids. Only the major part of the disordered fluorine atom is shown.

**Figure 2**

A portion of the crystal packing viewed along [010].

#### 4-[(1*E*)-3-(2,6-Dichloro-3-fluorophenyl)-3-oxoprop-1-en-1-yl]benzonitrile

##### Crystal data

$C_{16}H_8Cl_2FNO$

$M_r = 320.13$

Monoclinic,  $P2_1/c$

Hall symbol:  $-P\ 2_1/c$

$a = 13.2751$  (3) Å

$b = 8.5002$  (2) Å

$c = 13.9854$  (3) Å

$\beta = 116.773$  (1)°

$V = 1408.95$  (6) Å<sup>3</sup>

$Z = 4$

$F(000) = 648$

$D_x = 1.509$  Mg m<sup>-3</sup>

Melting point = 414–417 K

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 7660 reflections

$\theta = 2.9$ – $28.3$ °

$\mu = 0.47$  mm<sup>-1</sup>

$T = 200$  K

Platelet, green

$0.55 \times 0.32 \times 0.20$  mm

Data collection

Bruker APEXII CCD  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\varphi$  and  $\omega$  scans  
Absorption correction: multi-scan  
(SADABS; Bruker, 2008)  
 $T_{\min} = 0.783$ ,  $T_{\max} = 0.913$

12885 measured reflections  
3480 independent reflections  
2904 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.013$   
 $\theta_{\max} = 28.3^\circ$ ,  $\theta_{\min} = 2.9^\circ$   
 $h = -17 \rightarrow 17$   
 $k = -8 \rightarrow 11$   
 $l = -18 \rightarrow 18$

Refinement

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.037$   
 $wR(F^2) = 0.098$   
 $S = 1.05$   
3480 reflections  
200 parameters  
0 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0358P)^2 + 0.6287P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.41 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.43 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C11	0.32198 (4)	0.25664 (5)	0.03764 (4)	0.05826 (14)	
C12	0.43426 (5)	0.87196 (6)	0.08058 (4)	0.06998 (17)	
O1	0.23229 (11)	0.62352 (16)	-0.09939 (9)	0.0582 (3)	
N1	-0.11644 (13)	0.5937 (3)	0.40221 (14)	0.0785 (6)	
C1	0.21695 (12)	0.57997 (17)	0.14653 (12)	0.0392 (3)	
H1	0.2927	0.5473	0.1891	0.047*	
C2	0.18838 (14)	0.6169 (2)	0.04481 (13)	0.0460 (4)	
H2	0.1140	0.6544	0.0017	0.055*	
C3	0.26474 (14)	0.60326 (19)	-0.00432 (12)	0.0435 (3)	
C4	-0.06074 (13)	0.5902 (2)	0.35969 (14)	0.0525 (4)	
C11	0.14352 (12)	0.58457 (17)	0.19935 (11)	0.0370 (3)	
C12	0.03215 (13)	0.6397 (2)	0.14813 (13)	0.0487 (4)	
H12	0.0022	0.6771	0.0765	0.058*	
C13	-0.03410 (13)	0.6403 (2)	0.20037 (13)	0.0489 (4)	
H13	-0.1097	0.6771	0.1647	0.059*	
C14	0.00933 (12)	0.58722 (18)	0.30543 (12)	0.0408 (3)	
C15	0.11978 (12)	0.5340 (2)	0.35809 (12)	0.0440 (3)	
H15	0.1498	0.4984	0.4301	0.053*	
C16	0.18571 (12)	0.53320 (19)	0.30467 (12)	0.0407 (3)	
H16	0.2614	0.4968	0.3407	0.049*	
C21	0.38739 (13)	0.56076 (18)	0.06658 (11)	0.0394 (3)	
C22	0.42114 (12)	0.40474 (19)	0.08897 (11)	0.0404 (3)	
C24	0.61343 (14)	0.4808 (3)	0.19589 (13)	0.0538 (4)	
H24	0.6899	0.4539	0.2409	0.065*	
C26	0.47029 (15)	0.6754 (2)	0.10883 (13)	0.0467 (4)	

C23	0.53293 (14)	0.3657 (2)	0.15222 (13)	0.0474 (4)	
H23	0.5542	0.2582	0.1656	0.057*	0.470 (3)
F1B	0.6550 (2)	0.7502 (4)	0.2076 (2)	0.0847 (11)	0.470 (3)
C25	0.58158 (15)	0.6354 (2)	0.17347 (14)	0.0545 (4)	
H25	0.6367	0.7160	0.2027	0.065*	0.530 (3)
F1A	0.56085 (16)	0.2148 (3)	0.16948 (16)	0.0610 (7)	0.530 (3)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C11	0.0520 (2)	0.0417 (2)	0.0616 (3)	-0.00337 (17)	0.00832 (19)	0.00136 (18)
C12	0.1033 (4)	0.0436 (2)	0.0648 (3)	-0.0126 (2)	0.0394 (3)	-0.0003 (2)
O1	0.0635 (7)	0.0718 (9)	0.0403 (6)	0.0168 (6)	0.0242 (6)	0.0145 (6)
N1	0.0455 (8)	0.1321 (18)	0.0629 (10)	0.0129 (10)	0.0288 (8)	-0.0008 (11)
C1	0.0376 (7)	0.0386 (7)	0.0407 (7)	0.0002 (6)	0.0171 (6)	0.0005 (6)
C2	0.0444 (8)	0.0502 (9)	0.0432 (8)	0.0089 (7)	0.0195 (7)	0.0076 (7)
C3	0.0508 (8)	0.0405 (8)	0.0406 (8)	0.0047 (7)	0.0217 (7)	0.0052 (6)
C4	0.0361 (7)	0.0725 (12)	0.0473 (9)	0.0051 (8)	0.0172 (7)	-0.0027 (8)
C11	0.0355 (7)	0.0349 (7)	0.0389 (7)	-0.0007 (6)	0.0153 (6)	-0.0010 (6)
C12	0.0397 (8)	0.0596 (10)	0.0418 (8)	0.0076 (7)	0.0139 (6)	0.0099 (7)
C13	0.0336 (7)	0.0586 (10)	0.0497 (9)	0.0088 (7)	0.0146 (6)	0.0058 (7)
C14	0.0346 (7)	0.0428 (8)	0.0451 (8)	-0.0006 (6)	0.0182 (6)	-0.0052 (6)
C15	0.0392 (7)	0.0525 (9)	0.0404 (7)	0.0054 (6)	0.0179 (6)	0.0025 (7)
C16	0.0332 (7)	0.0465 (8)	0.0401 (7)	0.0059 (6)	0.0147 (6)	0.0024 (6)
C21	0.0461 (8)	0.0440 (8)	0.0327 (7)	0.0000 (6)	0.0219 (6)	0.0015 (6)
C22	0.0405 (7)	0.0443 (8)	0.0348 (7)	-0.0011 (6)	0.0155 (6)	-0.0012 (6)
C24	0.0390 (8)	0.0831 (13)	0.0393 (8)	-0.0030 (8)	0.0176 (7)	-0.0014 (8)
C26	0.0603 (9)	0.0472 (9)	0.0411 (8)	-0.0065 (7)	0.0303 (7)	-0.0005 (7)
C23	0.0464 (8)	0.0571 (10)	0.0389 (7)	0.0080 (7)	0.0193 (7)	0.0015 (7)
F1B	0.0738 (17)	0.107 (2)	0.0622 (16)	-0.0519 (16)	0.0211 (13)	-0.0060 (14)
C25	0.0535 (9)	0.0717 (12)	0.0433 (8)	-0.0211 (9)	0.0260 (8)	-0.0071 (8)
F1A	0.0529 (11)	0.0588 (13)	0.0594 (12)	0.0184 (9)	0.0147 (9)	0.0096 (9)

*Geometric parameters (Å, °)*

C11—C22	1.7267 (16)	C13—H13	0.9500
C12—C26	1.7338 (18)	C14—C15	1.387 (2)
O1—C3	1.2112 (18)	C15—C16	1.383 (2)
N1—C4	1.139 (2)	C15—H15	0.9500
C1—C2	1.334 (2)	C16—H16	0.9500
C1—C11	1.465 (2)	C21—C26	1.387 (2)
C1—H1	0.9500	C21—C22	1.390 (2)
C2—C3	1.463 (2)	C22—C23	1.383 (2)
C2—H2	0.9500	C24—C23	1.373 (3)
C3—C21	1.520 (2)	C24—C25	1.373 (3)
C4—C14	1.441 (2)	C24—H24	0.9500
C11—C16	1.390 (2)	C26—C25	1.383 (3)
C11—C12	1.402 (2)	C23—F1A	1.327 (3)

C12—C13	1.373 (2)	C23—H23	0.9500
C12—H12	0.9500	F1B—C25	1.308 (3)
C13—C14	1.390 (2)	C25—H25	0.9500
C2—C1—C11	126.67 (14)	C15—C16—H16	119.3
C2—C1—H1	116.7	C11—C16—H16	119.3
C11—C1—H1	116.7	C26—C21—C22	117.46 (14)
C1—C2—C3	123.90 (14)	C26—C21—C3	121.47 (15)
C1—C2—H2	118.1	C22—C21—C3	121.05 (14)
C3—C2—H2	118.1	C23—C22—C21	121.15 (15)
O1—C3—C2	121.68 (15)	C23—C22—C11	119.24 (13)
O1—C3—C21	119.79 (14)	C21—C22—C11	119.61 (11)
C2—C3—C21	118.53 (13)	C23—C24—C25	118.87 (16)
N1—C4—C14	179.4 (2)	C23—C24—H24	120.6
C16—C11—C12	118.39 (14)	C25—C24—H24	120.6
C16—C11—C1	118.96 (13)	C25—C26—C21	120.98 (17)
C12—C11—C1	122.65 (13)	C25—C26—C12	119.32 (14)
C13—C12—C11	120.68 (15)	C21—C26—C12	119.70 (13)
C13—C12—H12	119.7	F1A—C23—C24	120.69 (17)
C11—C12—H12	119.7	F1A—C23—C22	118.67 (17)
C12—C13—C14	120.06 (14)	C24—C23—C22	120.64 (17)
C12—C13—H13	120.0	F1A—C23—H23	1.0
C14—C13—H13	120.0	C24—C23—H23	119.7
C15—C14—C13	120.21 (14)	C22—C23—H23	119.7
C15—C14—C4	120.30 (14)	F1B—C25—C24	121.9 (2)
C13—C14—C4	119.48 (14)	F1B—C25—C26	117.1 (2)
C16—C15—C14	119.30 (14)	C24—C25—C26	120.87 (16)
C16—C15—H15	120.4	F1B—C25—H25	3.7
C14—C15—H15	120.4	C24—C25—H25	119.6
C15—C16—C11	121.34 (13)	C26—C25—H25	119.6
C11—C1—C2—C3	-177.25 (15)	C26—C21—C22—C23	0.9 (2)
C1—C2—C3—O1	172.76 (17)	C3—C21—C22—C23	179.07 (13)
C1—C2—C3—C21	-6.6 (2)	C26—C21—C22—C11	-179.43 (11)
C2—C1—C11—C16	176.31 (16)	C3—C21—C22—C11	-1.24 (19)
C2—C1—C11—C12	-4.1 (3)	C22—C21—C26—C25	-2.0 (2)
C16—C11—C12—C13	-1.1 (2)	C3—C21—C26—C25	179.84 (14)
C1—C11—C12—C13	179.29 (16)	C22—C21—C26—C12	178.12 (11)
C11—C12—C13—C14	0.6 (3)	C3—C21—C26—C12	-0.07 (19)
C12—C13—C14—C15	0.2 (3)	C25—C24—C23—F1A	178.08 (18)
C12—C13—C14—C4	179.32 (17)	C25—C24—C23—C22	-1.7 (2)
N1—C4—C14—C15	136 (21)	C21—C22—C23—F1A	-178.81 (16)
N1—C4—C14—C13	-43 (21)	C11—C22—C23—F1A	1.5 (2)
C13—C14—C15—C16	-0.5 (2)	C21—C22—C23—C24	0.9 (2)
C4—C14—C15—C16	-179.57 (16)	C11—C22—C23—C24	-178.75 (12)
C14—C15—C16—C11	0.0 (2)	C23—C24—C25—F1B	-176.2 (2)
C12—C11—C16—C15	0.8 (2)	C23—C24—C25—C26	0.6 (2)
C1—C11—C16—C15	-179.54 (14)	C21—C26—C25—F1B	178.18 (19)

O1—C3—C21—C26	84.6 (2)	C12—C26—C25—F1B	-1.9 (2)
C2—C3—C21—C26	-96.08 (18)	C21—C26—C25—C24	1.3 (2)
O1—C3—C21—C22	-93.55 (19)	C12—C26—C25—C24	-178.81 (13)
C2—C3—C21—C22	85.81 (19)		

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
C1—H1 $\cdots$ F1 <i>A</i> <sup>i</sup>	0.95	2.50	3.130 (2)	124

Symmetry code: (i)  $-x+1, y+1/2, -z+1/2$ .