

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1-(2-Fluorophenyl)-3-(2,4,6-trimethoxyphenyl)prop-2-en-1-one

M. Prabuswamy,^a S. Madan Kumar,^a D. Bhuvaneshwar,^b Ch. S. S. Murthy^b and N. K. Lokanath^{a*}

^aDepartment of Studies in Physics, Manasagangotri, University of Mysore, Mysore 570 006, India, and ^bDepartment of Nanotechnology, School of Interdisciplinary Courses, Noorul Islam Centre for Higher Education, Kumarcoil, Kanyakumari 629 180, India

Correspondence e-mail: lokanath@physics.uni-mysore.ac.in


Received 28 September 2012; accepted 16 October 2012

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.039; wR factor = 0.100; data-to-parameter ratio = 14.4.

In the title compound, $\text{C}_{18}\text{H}_{17}\text{FO}_4$, the dihedral angle between the aromatic rings is 32.29 (8°). The C atoms of the methoxy groups deviate from their attached ring plane by 0.018 (2), -0.006 (2) and -0.094 (2) Å. In the crystal, $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into $C(6)$ [001] chains.

Related literature

For the synthesis and properties of the title compound, see: Rimal *et al.* (2012). For a related structure, see: Jasinski *et al.* (2009).



Experimental

Crystal data

 $\text{C}_{18}\text{H}_{17}\text{FO}_4$ $M_r = 316.32$

Monoclinic, $P2_1/n$
 $a = 7.0927$ (3) Å
 $b = 25.9711$ (11) Å
 $c = 8.7487$ (4) Å
 $\beta = 91.584$ (4) $^\circ$
 $V = 1610.94$ (12) Å³

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 296$ K
 $0.22 \times 0.22 \times 0.20$ mm

Data collection

Oxford Diffraction Xcalibur Eos
 CCD diffractometer
 16217 measured reflections

3057 independent reflections
 1946 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.037$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.100$
 $S = 1.00$
 3057 reflections

212 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.17$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.14$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}2-\text{H}2\cdots\text{O}1^i$	0.93	2.44	3.338 (2)	162

Symmetry code: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2009); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *SHELXL97* and *Mercury*.

SMK thanks the UGC–BRS and the University of Mysore for the award of a fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6965).

References

- Jasinski, J. P., Butcher, R. J., Veena, K., Narayana, B. & Yathirajan, H. S. (2009). *Acta Cryst.* **E65**, o1965–o1966.
 Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
 Oxford Diffraction (2009). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, England.
 Rimal, I., SriBindu, M. C., Sudhakar, P. R., JosephJoly, V. L., Wu, R. J. & ChMurthy, S. S. S. (2012). *Adv. Sci. Eng. Med.* **4**, 1–7.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2012). E68, o3190 [doi:10.1107/S1600536812043139]

1-(2-Fluorophenyl)-3-(2,4,6-trimethoxyphenyl)prop-2-en-1-one

M. Prabuswamy, S. Madan Kumar, D. Bhuvaneshwar, Ch. S. S. S. Murthy and N. K. Lokanath

S1. Comment

As part of our ongoing studies of chalcones with possible biological activity (Rimal *et al.*, 2012), we now describe the crystal structure of 1-(2-fluorophenyl)-3-(2,4,6-trimethoxyphenyl)prop-2-en-1-one, (I).

The *ORTEP* drawing of the title molecule is as shown in Fig. 1. The dihedral angle between the aromatic rings of 2-fluorophenyl and trimethoxy phenyl group is 32.29 (8)°. Mean plane of prop-2-ene-1 one moiety makes an angle of 37.65 (9)° with 2-fluorophenyl and 11.64 (9)° with trimethoxyphenyl moiety. The overall geometry of the title compound is similar to that of (2E)-1-(4-fluorophenyl)-3-(3,4,5-trimethoxyphenyl)prop-2-en-1-one (Jasinski *et al.*, 2009).

The molecules are connected by C—H···O interactions (Fig. 2).

S2. Experimental

The title compound was synthesized as per the procedure reported in the literature (Rimal *et al.*, 2012). The final product was obtained by recrystallization using aqueous ethyl acetate: methanol as a solvent. Slow evaporation method yielded brown blocks.

S3. Refinement

All the hydrogen atoms of the compound are fixed geometrically (C—H = 0.93–0.97 Å) and allowed to ride on their parent atoms.





Figure 1

ORTEP diagram of the title compound with 50% probability ellipsoids.

**Figure 2**

Packing diagram of the title compound, viewed along the crystallographic *a* axis. C—H...O hydrogen bonds are indicated by dashed lines.

1-(2-Fluorophenyl)-3-(2,4,6-trimethoxyphenyl)prop-2-en-1-one

Crystal data

$C_{18}H_{17}FO_4$

$M_r = 316.32$

Monoclinic, $P2_1/n$

Hall symbol: $-P\ 2_1/n$

$a = 7.0927$ (3) Å

$b = 25.9711$ (11) Å

$c = 8.7487$ (4) Å

$\beta = 91.584$ (4)°

$V = 1610.94$ (12) Å³

$Z = 4$

$F(000) = 664$

$D_x = 1.304$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3057 reflections

$\theta = 2.5$ – 25.7 °

$\mu = 0.10$ mm⁻¹

$T = 296$ K

Block, brown

$0.22 \times 0.22 \times 0.20$ mm

Data collection

Oxford Diffraction Xcalibur Eos CCD diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 16.0839 pixels mm⁻¹

ω scans

16217 measured reflections

3057 independent reflections

1946 reflections with $I > 2\sigma(I)$

$R_{int} = 0.037$

$\theta_{max} = 25.7$ °, $\theta_{min} = 2.5$ °

$h = -8 \rightarrow 8$

$k = -31 \rightarrow 30$

$l = -10 \rightarrow 10$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.039$

$wR(F^2) = 0.100$

$S = 1.00$

3057 reflections

212 parameters

0 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0497P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{max} < 0.001$

$\Delta\rho_{max} = 0.17$ e Å⁻³

$\Delta\rho_{min} = -0.14$ e Å⁻³

Extinction correction: *SHELXL97* (Sheldrick, 2008), $FC^* = KFC[1 + 0.001XFC^2\Lambda^3/SIN(2\Theta)]^{-1/4}$

Extinction coefficient: 0.023 (2)

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement on F^2 for ALL reflections except those flagged by the user for potential systematic errors. Weighted R -factors wR and all goodnesses of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The observed criterion of $F^2 > \sigma(F^2)$ is used only for calculating $-R$ -factor-obs *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
F1	0.47580 (14)	0.21236 (4)	0.09672 (13)	0.0862 (5)
O1	0.05893 (17)	0.14446 (4)	0.32501 (13)	0.0687 (4)
O2	0.65872 (14)	0.07225 (4)	0.09354 (13)	0.0565 (4)
O3	0.81263 (17)	-0.10017 (4)	0.20765 (13)	0.0681 (5)
O4	0.24772 (16)	-0.02139 (4)	0.39762 (13)	0.0641 (4)
C1	0.2939 (2)	0.22710 (6)	0.06805 (19)	0.0569 (7)
C2	0.2679 (3)	0.27123 (7)	-0.0164 (2)	0.0709 (8)
C3	0.0855 (4)	0.28686 (7)	-0.0490 (2)	0.0775 (9)
C4	-0.0645 (3)	0.25900 (7)	0.0031 (2)	0.0724 (8)
C5	-0.0318 (2)	0.21601 (6)	0.09185 (19)	0.0583 (6)
C6	0.1499 (2)	0.19858 (6)	0.12600 (17)	0.0463 (5)
C7	0.1721 (2)	0.15104 (6)	0.22354 (17)	0.0464 (5)
C8	0.3191 (2)	0.11421 (6)	0.19021 (17)	0.0471 (5)
C9	0.3266 (2)	0.06858 (5)	0.26294 (17)	0.0444 (5)
C10	0.4532 (2)	0.02558 (5)	0.24723 (16)	0.0422 (5)
C11	0.6197 (2)	0.02680 (6)	0.16292 (16)	0.0448 (5)
C12	0.7362 (2)	-0.01548 (6)	0.15323 (17)	0.0516 (6)
C13	0.6887 (2)	-0.06078 (6)	0.22712 (18)	0.0510 (6)
C14	0.5274 (2)	-0.06425 (6)	0.31166 (18)	0.0519 (6)
C15	0.4117 (2)	-0.02141 (6)	0.32072 (16)	0.0463 (6)
C16	0.8256 (2)	0.07605 (6)	0.00797 (19)	0.0596 (7)
C17	0.7766 (3)	-0.14799 (6)	0.2812 (2)	0.0782 (8)
C18	0.1869 (3)	-0.06735 (6)	0.4707 (2)	0.0699 (7)
H2	0.37020	0.29000	-0.05050	0.0850*
H3	0.06380	0.31650	-0.10660	0.0930*
H4	-0.18730	0.26920	-0.02150	0.0870*
H5	-0.13400	0.19810	0.13000	0.0700*
H8	0.40860	0.12220	0.11820	0.0570*
H9	0.23470	0.06400	0.33550	0.0530*
H12	0.84580	-0.01370	0.09760	0.0620*
H14	0.49710	-0.09470	0.36140	0.0620*
H16A	0.93360	0.07030	0.07430	0.0890*
H16B	0.83310	0.10980	-0.03650	0.0890*
H16C	0.82270	0.05060	-0.07170	0.0890*
H17A	0.77770	-0.14300	0.39000	0.1170*

H17B	0.87250	-0.17240	0.25570	0.1170*
H17C	0.65550	-0.16080	0.24760	0.1170*
H18A	0.16980	-0.09410	0.39580	0.1050*
H18B	0.06970	-0.06100	0.51960	0.1050*
H18C	0.28040	-0.07790	0.54580	0.1050*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
F1	0.0529 (7)	0.1002 (8)	0.1054 (9)	-0.0147 (6)	0.0030 (6)	0.0333 (7)
O1	0.0699 (8)	0.0614 (7)	0.0766 (8)	0.0155 (6)	0.0370 (7)	0.0142 (6)
O2	0.0464 (7)	0.0508 (7)	0.0734 (8)	0.0010 (5)	0.0218 (6)	0.0077 (6)
O3	0.0714 (9)	0.0548 (7)	0.0788 (8)	0.0200 (6)	0.0136 (7)	0.0074 (6)
O4	0.0553 (8)	0.0594 (7)	0.0789 (8)	-0.0048 (6)	0.0233 (6)	0.0108 (6)
C1	0.0501 (12)	0.0633 (11)	0.0573 (11)	-0.0054 (9)	0.0000 (9)	0.0046 (9)
C2	0.0886 (16)	0.0605 (12)	0.0639 (12)	-0.0176 (11)	0.0057 (11)	0.0126 (10)
C3	0.1100 (19)	0.0544 (12)	0.0676 (13)	0.0113 (13)	-0.0069 (13)	0.0107 (10)
C4	0.0753 (15)	0.0611 (12)	0.0803 (14)	0.0190 (11)	-0.0039 (11)	0.0038 (10)
C5	0.0552 (12)	0.0542 (10)	0.0659 (11)	0.0074 (9)	0.0078 (9)	-0.0036 (9)
C6	0.0465 (10)	0.0444 (9)	0.0482 (9)	-0.0005 (8)	0.0055 (8)	-0.0024 (7)
C7	0.0435 (9)	0.0471 (9)	0.0490 (9)	-0.0032 (8)	0.0067 (8)	-0.0024 (7)
C8	0.0405 (9)	0.0525 (10)	0.0488 (9)	0.0013 (8)	0.0083 (7)	0.0004 (8)
C9	0.0383 (9)	0.0505 (9)	0.0447 (9)	-0.0028 (7)	0.0059 (7)	-0.0032 (7)
C10	0.0369 (9)	0.0464 (9)	0.0435 (8)	-0.0006 (7)	0.0024 (7)	-0.0016 (7)
C11	0.0427 (9)	0.0456 (9)	0.0462 (9)	-0.0009 (8)	0.0021 (7)	-0.0007 (7)
C12	0.0444 (10)	0.0573 (10)	0.0536 (10)	0.0035 (8)	0.0095 (8)	-0.0001 (8)
C13	0.0515 (11)	0.0504 (10)	0.0510 (10)	0.0083 (8)	-0.0013 (8)	-0.0044 (8)
C14	0.0576 (11)	0.0445 (9)	0.0535 (10)	-0.0025 (8)	0.0013 (9)	0.0026 (8)
C15	0.0419 (10)	0.0514 (10)	0.0457 (9)	-0.0057 (8)	0.0045 (8)	-0.0005 (7)
C16	0.0488 (11)	0.0609 (11)	0.0700 (12)	-0.0068 (8)	0.0182 (9)	-0.0010 (9)
C17	0.0957 (16)	0.0518 (11)	0.0871 (14)	0.0165 (11)	0.0049 (12)	0.0052 (10)
C18	0.0680 (13)	0.0693 (12)	0.0732 (13)	-0.0206 (10)	0.0182 (10)	0.0127 (10)

Geometric parameters (Å, °)

F1—C1	1.3624 (18)	C12—C13	1.388 (2)
O1—C7	1.2251 (19)	C13—C14	1.382 (2)
O2—C11	1.3593 (19)	C14—C15	1.386 (2)
O2—C16	1.4213 (18)	C2—H2	0.9300
O3—C13	1.3625 (19)	C3—H3	0.9300
O3—C17	1.4251 (19)	C4—H4	0.9300
O4—C15	1.3596 (18)	C5—H5	0.9300
O4—C18	1.427 (2)	C8—H8	0.9300
C1—C2	1.373 (2)	C9—H9	0.9300
C1—C6	1.370 (2)	C12—H12	0.9300
C2—C3	1.378 (3)	C14—H14	0.9300
C3—C4	1.375 (3)	C16—H16A	0.9600
C4—C5	1.376 (2)	C16—H16B	0.9600

C5—C6	1.391 (2)	C16—H16C	0.9600
C6—C7	1.507 (2)	C17—H17A	0.9600
C7—C8	1.451 (2)	C17—H17B	0.9600
C8—C9	1.345 (2)	C17—H17C	0.9600
C9—C10	1.4419 (19)	C18—H18A	0.9600
C10—C11	1.410 (2)	C18—H18B	0.9600
C10—C15	1.414 (2)	C18—H18C	0.9600
C11—C12	1.378 (2)		
C11—O2—C16	118.58 (11)	C3—C2—H2	121.00
C13—O3—C17	118.26 (13)	C2—C3—H3	120.00
C15—O4—C18	119.66 (13)	C4—C3—H3	120.00
F1—C1—C2	116.49 (15)	C3—C4—H4	120.00
F1—C1—C6	119.46 (14)	C5—C4—H4	120.00
C2—C1—C6	124.04 (15)	C4—C5—H5	119.00
C1—C2—C3	117.94 (18)	C6—C5—H5	119.00
C2—C3—C4	120.46 (17)	C7—C8—H8	120.00
C3—C4—C5	119.62 (19)	C9—C8—H8	120.00
C4—C5—C6	121.75 (15)	C8—C9—H9	115.00
C1—C6—C5	116.13 (14)	C10—C9—H9	115.00
C1—C6—C7	125.82 (13)	C11—C12—H12	120.00
C5—C6—C7	118.05 (13)	C13—C12—H12	120.00
O1—C7—C6	117.67 (13)	C13—C14—H14	121.00
O1—C7—C8	122.87 (14)	C15—C14—H14	121.00
C6—C7—C8	119.41 (13)	O2—C16—H16A	109.00
C7—C8—C9	120.33 (13)	O2—C16—H16B	109.00
C8—C9—C10	130.82 (14)	O2—C16—H16C	109.00
C9—C10—C11	124.42 (13)	H16A—C16—H16B	110.00
C9—C10—C15	119.15 (13)	H16A—C16—H16C	109.00
C11—C10—C15	116.43 (13)	H16B—C16—H16C	110.00
O2—C11—C10	115.91 (13)	O3—C17—H17A	109.00
O2—C11—C12	122.32 (13)	O3—C17—H17B	109.00
C10—C11—C12	121.77 (14)	O3—C17—H17C	110.00
C11—C12—C13	119.53 (13)	H17A—C17—H17B	109.00
O3—C13—C12	114.35 (13)	H17A—C17—H17C	109.00
O3—C13—C14	124.35 (14)	H17B—C17—H17C	109.00
C12—C13—C14	121.30 (14)	O4—C18—H18A	109.00
C13—C14—C15	118.58 (14)	O4—C18—H18B	109.00
O4—C15—C10	114.60 (13)	O4—C18—H18C	109.00
O4—C15—C14	122.99 (13)	H18A—C18—H18B	110.00
C10—C15—C14	122.39 (13)	H18A—C18—H18C	109.00
C1—C2—H2	121.00	H18B—C18—H18C	110.00
C16—O2—C11—C10	179.43 (13)	O1—C7—C8—C9	5.6 (2)
C16—O2—C11—C12	0.1 (2)	C6—C7—C8—C9	-171.60 (14)
C17—O3—C13—C12	179.14 (14)	C7—C8—C9—C10	178.00 (14)
C17—O3—C13—C14	-1.2 (2)	C8—C9—C10—C11	10.4 (3)
C18—O4—C15—C10	177.28 (13)	C8—C9—C10—C15	-169.93 (15)

C18—O4—C15—C14	-1.3 (2)	C9—C10—C11—O2	0.1 (2)
F1—C1—C2—C3	179.23 (15)	C9—C10—C11—C12	179.39 (14)
C6—C1—C2—C3	-2.2 (3)	C15—C10—C11—O2	-179.62 (12)
F1—C1—C6—C5	-179.97 (13)	C15—C10—C11—C12	-0.3 (2)
F1—C1—C6—C7	0.8 (2)	C9—C10—C15—O4	1.86 (19)
C2—C1—C6—C5	1.5 (2)	C9—C10—C15—C14	-179.54 (14)
C2—C1—C6—C7	-177.78 (16)	C11—C10—C15—O4	-178.45 (12)
C1—C2—C3—C4	0.5 (3)	C11—C10—C15—C14	0.2 (2)
C2—C3—C4—C5	1.8 (3)	O2—C11—C12—C13	179.81 (13)
C3—C4—C5—C6	-2.5 (3)	C10—C11—C12—C13	0.5 (2)
C4—C5—C6—C1	0.9 (2)	C11—C12—C13—O3	179.05 (13)
C4—C5—C6—C7	-179.77 (15)	C11—C12—C13—C14	-0.6 (2)
C1—C6—C7—O1	144.10 (16)	O3—C13—C14—C15	-179.15 (14)
C1—C6—C7—C8	-38.5 (2)	C12—C13—C14—C15	0.5 (2)
C5—C6—C7—O1	-35.2 (2)	C13—C14—C15—O4	178.24 (14)
C5—C6—C7—C8	142.22 (15)	C13—C14—C15—C10	-0.2 (2)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C2—H2...O1 ⁱ	0.93	2.44	3.338 (2)	162

Symmetry code: (i) $x+1/2, -y+1/2, z-1/2$.