

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4-(Methylsulfonyl)piperazin-1-ium chloride

 Hoong-Kun Fun,^{a,*}‡ Chin Sing Yeap,^{a,§} C. S. Chidan Kumar,^b H. S. Yathirajan^b and B. Narayana^c

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, ^bDepartment of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore 570 006, India, and ^cDepartment of Studies in Chemistry, Mangalore University, Mangalagangotri 574 199, India
Correspondence e-mail: hkfun@usm.my


Received 10 January 2010; accepted 10 January 2010

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.001$ Å; R factor = 0.023; wR factor = 0.072; data-to-parameter ratio = 32.1.

In the title molecular salt, $\text{C}_5\text{H}_{13}\text{N}_2\text{O}_2\text{S}^+\cdot\text{Cl}^-$, the complete cation is generated by crystallographic mirror symmetry, with both N atoms, the S atom and one C atom lying on the reflecting plane. The chloride ion also lies on the mirror plane. The piperazinium ring adopts a chair conformation and the N—S bond adopts an equatorial orientation. In the crystal structure, the component ions are linked into a three-dimensional framework by intermolecular N—H \cdots Cl and C—H \cdots Cl hydrogen bonds.

Related literature

For medicinal background to piperazine derivatives, see: Dinsmore & Beshore (2002); Berkheij *et al.* (2005); Humle & Cherrier (1999). For related structures, see: Bart *et al.* (1978); Girisha *et al.* (2008); Homrighausen & Krause Bauer (2002); Jin *et al.* (2007); Kubo *et al.* (2007); Parkin *et al.* (2004); Shen *et al.* (2006), Wang *et al.* (2006). For ring conformations, see: Cremer & Pople (1975). For the stability of the temperature controller used for the data collection, see: Cosier & Glazer (1986).



‡ Thomson Reuters ResearcherID: A-3561-2009.

§ Thomson Reuters ResearcherID: A-5523-2009.

Experimental

Crystal data

$\text{C}_5\text{H}_{13}\text{N}_2\text{O}_2\text{S}^+\cdot\text{Cl}^-$
 $M_r = 200.68$
 Monoclinic, $P2_1/m$
 $a = 6.0231$ (1) Å
 $b = 9.1097$ (2) Å
 $c = 7.9852$ (2) Å
 $\beta = 100.700$ (1)°
 $V = 430.52$ (2) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.64$ mm⁻¹
 $T = 100$ K
 $0.36 \times 0.32 \times 0.05$ mm

Data collection

Bruker APEX Duo CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2009)
 $T_{\min} = 0.801$, $T_{\max} = 0.968$
 10626 measured reflections
 2790 independent reflections
 2419 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.022$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.023$
 $wR(F^2) = 0.072$
 $S = 1.10$
 2790 reflections
 87 parameters
 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.49$ e Å⁻³
 $\Delta\rho_{\min} = -0.40$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1N1}\cdots\text{Cl}^{\text{i}}$	0.92 (2)	2.40 (2)	3.1341 (8)	137 (1)
$\text{N1}-\text{H2N1}\cdots\text{Cl}^{\text{ii}}$	0.93 (2)	2.19 (2)	3.0966 (8)	164 (1)
$\text{C1}-\text{H1A}\cdots\text{Cl}^{\text{iii}}$	0.953 (12)	2.700 (12)	3.5251 (6)	145.2 (9)
$\text{C3}-\text{H3A}\cdots\text{Cl}^{\text{i}}$	0.94 (2)	2.65 (2)	3.5487 (10)	160 (2)

Symmetry codes: (i) $x + 1, y, z - 1$; (ii) $x, y, z - 1$; (iii) $-x + 1, y - \frac{1}{2}, -z + 1$.

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

HKF thanks Universiti Sains Malaysia (USM) for the Research University Golden Goose Grant (No. 1001/PFIZIK/811012). CSY thanks USM for the award of a USM Fellowship. CSC thanks University of Mysore for research facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5306).

References

- Bart, J. C. J., Bassi, I. W. & Scordamaglia, R. (1978). *Acta Cryst.* **B34**, 2760–2764.
 Berkheij, M., van der Sluis, L., Sewing, C., den Boer, D. J., Terpstra, J. W., Heimstra, H., Bakker, W. I. L., van den Hoogen Band, A. & van Maarseveen, J. H. (2005). *Tetrahedron*, **46**, 2369–2371.
 Bruker (2009). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
 Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
 Dinsmore, C. J. & Beshore, D. C. (2002). *Tetrahedron*, **58**, 3297–3312.
 Girisha, H. R., Naveen, S., Vinaya, K., Sridhar, M. A., Shashidhara Prasad, J. & Rangappa, K. S. (2008). *Acta Cryst.* **E64**, o358.
 Homrighausen, C. L. & Krause Bauer, J. A. (2002). *Acta Cryst.* **E58**, o1395–o1396.

- Humle, C. & Cherrier, M. P. (1999). *Tetrahedron Lett.* **40**, 5295–5299.
- Jin, H.-M., Li, P.-F., Li, C.-Y. & Liu, B. (2007). *Acta Cryst. E63*, o3689.
- Kubo, K., Yamamoto, E., Hayakawa, A., Sakurai, T. & Mori, A. (2007). *Acta Cryst. E63*, o1347–o1348.
- Parkin, A., Oswald, I. D. H. & Parsons, S. (2004). *Acta Cryst. B60*, 219–227.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Shen, L., Wang, F.-W., Cheng, A.-B. & Yang, S. (2006). *Acta Cryst. E62*, o2242–o2244.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
- Wang, J., Zeng, T., Li, M.-L., Duan, E.-H. & Li, J.-S. (2006). *Acta Cryst. E62*, o2912–o2913.

supporting information

Acta Cryst. (2010). E66, o361–o362 [https://doi.org/10.1107/S1600536810001224]

4-(Methylsulfonyl)piperazin-1-ium chloride

Hoong-Kun Fun, Chin Sing Yeap, C. S. Chidan Kumar, H. S. Yathirajan and B. Narayana

S1. Comment

Piperazines are among the most important building blocks in today's drug discovery. The piperazine nucleus is capable of binding to multiple receptors with high affinity and therefore piperazine has been classified as a privileged structure (Dinsmore & Beshore, 2002). They are found in biologically active compounds across a number of different therapeutic areas (Berkheij *et al.*, 2005) such as antifungal, antibacterial, antimalarial, antipsychotic, antidepressant and antitumour activity against colon, prostate, breast, lung and leukemia tumors (Humble & Cherrier, 1999). The piperazines are a broad class of chemical compounds, many with important pharmacological properties, which contain a core piperazine functional group. 1-(Methylsulfonyl)piperazine is an important intermediate in synthetic organic chemistry, mainly used as a pharmaceutical intermediate.

The crystal structures of *trans*-2,5-dimethylpiperazine dihydrochloride (Bart *et al.*, 1978), 1-(3-chlorophenyl)-4-(3-chloropropyl)piperazinium chloride (Homrighausen & Krause Bauer, 2002), piperazine (Parkin *et al.*, 2004), 2,2'-(piperazine-1,4-dium-1,4-diyl)diacetate dehydrate (Shen *et al.*, 2006), 1,4-bis(chloroacetyl)piperazine (Wang *et al.*, 2006), 1,4-bis(1-naphthylmethyl) piperazine (Kubo *et al.*, 2007), 1,4-bis(4-chlorobenzo-yl)piperazine (Jin *et al.*, 2007) and 1-benzhydryl-4-(4-chlorophenylsulfonyl) piperazine (Girisha *et al.*, 2008) have been reported. In view of the importance of the title compound, this paper reports its crystal structure.

The asymmetric unit of the title compound contains one-half of a cation and half of a chloride anion (Fig. 1). The C11, S1, N1, N2, and C3 atoms are lying on a mirror plane. The piperazinium ring adopts a chair conformation with puckering amplitude $Q = 0.5680(7) \text{ \AA}$, $\theta = 179.90(7)^\circ$, $\varphi = 180(7)^\circ$ (Cremer & Pople, 1975). In the crystal structure (Fig. 2), the molecules are linked into a three-dimensional framework by intermolecular hydrogen bonds (Table 1).

S2. Experimental

The title compound was obtained as a gift sample from *R. L. Fine Chem.*, Bangalore, India. The compound was used without further purification. Colourless plates of (I) were obtained from slow evaporation of a methanol solution (m.p.: 489–492 K).

S3. Refinement

All H atoms were located in a difference Fourier map and refined freely.




Figure 1

The molecular structure of (I) with 50% probability ellipsoids for the non-H atoms. Atoms with suffix A are generated by the symmetry operation $(x, 1/2 - y, z)$.




Figure 2

The crystal packing of (I), viewed down the a axis, showing the hydrogen-bonded (dashed lines) three-dimensional framework.

4-(Methylsulfonyl)piperazin-1-ium chloride

Crystal data

$C_5H_{13}N_2O_2S^+Cl^-$

$M_r = 200.68$

Monoclinic, $P2_1/m$

Hall symbol: $-P\ 2yb$

$a = 6.0231$ (1) Å

$b = 9.1097$ (2) Å

$c = 7.9852$ (2) Å

$\beta = 100.700$ (1)°

$V = 430.52$ (2) Å³

$Z = 2$

$F(000) = 212$

$D_x = 1.548$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4890 reflections

$\theta = 3.4\text{--}40.1^\circ$

$\mu = 0.64$ mm⁻¹

$T = 100$ K

Plate, colourless

$0.36 \times 0.32 \times 0.05$ mm

Data collection

Bruker APEX Duo CCD diffractometer	10626 measured reflections
Radiation source: fine-focus sealed tube	2790 independent reflections
Graphite monochromator	2419 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.022$
Absorption correction: multi-scan (SADABS; Bruker, 2009)	$\theta_{\text{max}} = 40.1^\circ$, $\theta_{\text{min}} = 2.6^\circ$
$T_{\text{min}} = 0.801$, $T_{\text{max}} = 0.968$	$h = -10 \rightarrow 10$
	$k = -14 \rightarrow 16$
	$l = -14 \rightarrow 14$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.023$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.072$	$w = 1/[\sigma^2(F_o^2) + (0.0335P)^2 + 0.0922P]$
$S = 1.10$	where $P = (F_o^2 + 2F_c^2)/3$
2790 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
87 parameters	$\Delta\rho_{\text{max}} = 0.49 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.40 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cyrosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.30674 (3)	0.2500	0.93448 (3)	0.01195 (5)
S1	0.69091 (3)	0.2500	0.56598 (2)	0.00951 (5)
N1	0.82854 (12)	0.2500	0.03611 (9)	0.01007 (11)
N2	0.79320 (12)	0.2500	0.38830 (9)	0.00974 (11)
O1	0.75935 (9)	0.11396 (6)	0.65235 (6)	0.01498 (9)
C1	0.87784 (10)	0.11522 (7)	0.14202 (8)	0.01165 (9)
C2	0.74465 (10)	0.11465 (7)	0.28537 (8)	0.01168 (9)
C3	0.39411 (15)	0.2500	0.50705 (12)	0.01332 (13)
H1A	0.8351 (18)	0.0315 (13)	0.0719 (14)	0.012 (2)*
H1B	1.040 (2)	0.1186 (13)	0.1845 (16)	0.016 (3)*
H2A	0.583 (2)	0.1017 (14)	0.2371 (15)	0.018 (3)*
H2B	0.789 (2)	0.0335 (16)	0.3554 (17)	0.027 (3)*
H3A	0.331 (3)	0.2500	0.606 (2)	0.019 (4)*
H3B	0.350 (2)	0.3381 (15)	0.4460 (16)	0.025 (3)*

H1N1	0.914 (3)	0.2500	-0.048 (2)	0.019 (4)*
H2N1	0.677 (3)	0.2500	-0.017 (2)	0.021 (4)*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.00886 (7)	0.01261 (8)	0.01485 (9)	0.000	0.00346 (6)	0.000
S1	0.01110 (8)	0.01033 (8)	0.00696 (8)	0.000	0.00128 (6)	0.000
N1	0.0098 (2)	0.0115 (3)	0.0095 (3)	0.000	0.00316 (19)	0.000
N2	0.0126 (2)	0.0084 (2)	0.0087 (2)	0.000	0.0032 (2)	0.000
O1	0.01874 (19)	0.01499 (19)	0.01109 (18)	0.00356 (16)	0.00245 (15)	0.00480 (15)
C1	0.0148 (2)	0.00900 (19)	0.0122 (2)	0.00097 (17)	0.00547 (17)	-0.00041 (17)
C2	0.0162 (2)	0.0083 (2)	0.0117 (2)	-0.00109 (16)	0.00571 (17)	-0.00060 (16)
C3	0.0115 (3)	0.0166 (3)	0.0121 (3)	0.000	0.0027 (2)	0.000

Geometric parameters (Å, °)

S1—O1 ⁱ	1.4408 (5)	N2—C2 ⁱ	1.4806 (7)
S1—O1	1.4408 (5)	C1—C2	1.5148 (8)
S1—N2	1.6484 (7)	C1—H1A	0.953 (11)
S1—C3	1.7621 (9)	C1—H1B	0.976 (12)
N1—C1	1.4892 (7)	C2—H2A	0.983 (13)
N1—C1 ⁱ	1.4892 (7)	C2—H2B	0.935 (14)
N1—H1N1	0.920 (17)	C3—H3A	0.941 (18)
N1—H2N1	0.933 (19)	C3—H3B	0.951 (14)
N2—C2	1.4806 (7)		
O1 ⁱ —S1—O1	118.67 (4)	N1—C1—C2	110.75 (5)
O1 ⁱ —S1—N2	107.01 (2)	N1—C1—H1A	108.8 (7)
O1—S1—N2	107.01 (2)	C2—C1—H1A	108.5 (6)
O1 ⁱ —S1—C3	108.28 (3)	N1—C1—H1B	104.6 (7)
O1—S1—C3	108.29 (3)	C2—C1—H1B	112.1 (7)
N2—S1—C3	107.03 (4)	H1A—C1—H1B	112.0 (9)
C1—N1—C1 ⁱ	111.07 (7)	N2—C2—C1	109.81 (5)
C1—N1—H1N1	109.7 (5)	N2—C2—H2A	113.4 (7)
C1 ⁱ —N1—H1N1	109.7 (5)	C1—C2—H2A	109.1 (7)
C1—N1—H2N1	109.5 (5)	N2—C2—H2B	108.7 (8)
C1 ⁱ —N1—H2N1	109.5 (5)	C1—C2—H2B	108.7 (7)
H1N1—N1—H2N1	107.3 (15)	H2A—C2—H2B	107.1 (10)
C2—N2—C2 ⁱ	112.77 (7)	S1—C3—H3A	108.9 (11)
C2—N2—S1	114.27 (4)	S1—C3—H3B	108.1 (8)
C2 ⁱ —N2—S1	114.27 (4)	H3A—C3—H3B	108.3 (9)
O1 ⁱ —S1—N2—C2	178.10 (5)	C3—S1—N2—C2 ⁱ	66.00 (5)
O1—S1—N2—C2	49.91 (6)	C1 ⁱ —N1—C1—C2	56.95 (8)
C3—S1—N2—C2	-65.99 (5)	C2 ⁱ —N2—C2—C1	56.73 (8)

O1 ⁱ —S1—N2—C2 ⁱ	-49.91 (6)	S1—N2—C2—C1	-170.56 (4)
O1—S1—N2—C2 ⁱ	-178.10 (5)	N1—C1—C2—N2	-55.89 (7)

Symmetry code: (i) $x, -y+1/2, z$.

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
N1—H1M1...C11 ⁱⁱ	0.92 (2)	2.40 (2)	3.1341 (8)	137 (1)
N1—H2M1...C11 ⁱⁱⁱ	0.93 (2)	2.19 (2)	3.0966 (8)	164 (1)
C1—H1A...C11 ^{iv}	0.953 (12)	2.700 (12)	3.5251 (6)	145.2 (9)
C3—H3A...C11	0.94 (2)	2.65 (2)	3.5487 (10)	160 (2)

Symmetry codes: (ii) $x+1, y, z-1$; (iii) $x, y, z-1$; (iv) $-x+1, y-1/2, -z+1$.