

H. G. Anilkumar,^a
 H. S. Yathirajan,^a
 T. Narasimhamurthy,^b
 T. Vijay^c and R. S. Rathore^{d*}

^aDepartment of Studies in Chemistry, University of Mysore, Manasagangothri, Mysore 570 006, India, ^bBioinformatics Center, Indian Institute of Science, Bangalore 560 012, India, ^cSolid State and Structural Chemistry Unit, Indian Institute of Science, Bangalore 560 012, India, and ^dOriental Organization of Molecular and Structural Biology, 204 Agarwal Bhavan, Malleshwaram, Bangalore 560 055, India

Correspondence e-mail:
 ravindranath_rathore@yahoo.com

Key indicators

Single-crystal X-ray study
 $T = 293\text{ K}$
 Mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$
 R factor = 0.038
 wR factor = 0.114
 Data-to-parameter ratio = 11.1


For details of how these key indicators were automatically derived from the article, see <http://journals.iucr.org/e>.

Intermolecularly associated carboxylic acid dimers in the supramolecular assembly of 2-(2-phenylethyl)benzoic acid

The molecular structure of the title acid, $\text{C}_{15}\text{H}_{14}\text{O}_2$, exhibits a *trans*-planar arrangement of the aromatic rings. The carboxyl group is inclined at an angle of $6.4(1)^\circ$ with respect to the attached benzene ring. The classical hydrogen-bonded carboxylic acid dimers ($\text{O}\cdots\text{O} = 2.7\text{ \AA}$), characterized by an $R_2^2(8)$ pattern, predominantly stabilize the supramolecular assembly.

Comment

Benzoic acid was among the first organic crystal structures to be examined by the application of the X-ray crystallography (Bragg, 1921, 1922). Since then, numerous structures of benzoic acid and its derivatives have been reported (Sim *et al.*, 1955; Parkin *et al.*, 2005), owing to their importance in synthetic medicinal chemistry.



In the present report, we discuss the structure of one of these derivatives and the intermolecular interactions which govern its supramolecular assembly. The title acid, (I), is an intermediate in the preparation of dibenzosuberone (Ramesha & Roy, 2001) and the tricyclic antidepressant amitriptyline (Barbui *et al.*, 2004; Maizels & McCarberg, 2005).




Figure 1
 A view of (I), with the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms are shown as small spheres of arbitrary radii.

Received 6 July 2005
 Accepted 21 July 2005
 Online 27 July 2005

The molecule, shown in Fig. 1, assumes an overall *trans* conformation; $\omega(\text{C6}—\text{C7}—\text{C8}—\text{C9}) = 175.8 (1)^\circ$. The aromatic rings, linked by an ethylene bridge, are nearly parallel, with a dihedral angle of $4.3 (1)^\circ$. The plane of the carboxyl group is twisted by $6.4 (1)^\circ$ out of the plane of the attached benzene ring ($\text{C9}—\text{C14}$).

The characteristic feature of the molecular packing is the formation of classical hydrogen-bonded carboxylic acid dimers (Fig. 2). The $\text{O2}—\text{H2A} \cdots \text{O1}^i$ hydrogen-bond interaction involving centrosymmetrically related molecules is very strong (Table 1). The carboxylic acid dimers are characterized by an $R_2^2(8)$ ring pattern (Bernstein *et al.*, 1995). There are no significant overlaps of aromatic rings.

Experimental

The title acid was obtained from Arvee Pharma Mysore, India. Crystals were grown by slow evaporation of an ethanol solution.

Crystal data

$\text{C}_{15}\text{H}_{14}\text{O}_2$	$D_x = 1.227 \text{ Mg m}^{-3}$
$M_r = 226.26$	Mo $K\alpha$ radiation
Monoclinic, $P2_1/c$	Cell parameters from 1036 reflections
$a = 13.7005 (2) \text{ \AA}$	$\theta = 5–35^\circ$
$b = 6.0509 (1) \text{ \AA}$	$\mu = 0.08 \text{ mm}^{-1}$
$c = 15.1789 (1) \text{ \AA}$	$T = 293 (2) \text{ K}$
$\beta = 103.155 (1)^\circ$	Block, colorless
$V = 1225.32 (3) \text{ \AA}^3$	$0.14 \times 0.12 \times 0.08 \text{ mm}$
$Z = 4$	

Data collection

Siemens SMART CCD area-detector diffractometer	1421 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.024$
Absorption correction: none	$\theta_{\text{max}} = 23.3^\circ$
4864 measured reflections	$h = -15 \rightarrow 14$
1761 independent reflections	$k = -6 \rightarrow 6$
	$l = -16 \rightarrow 14$

Refinement

Refinement on F^2	$w = 1/[\sigma^2(F_o^2) + (0.0746P)^2]$
$R[F^2 > 2\sigma(F^2)] = 0.038$	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.114$	$(\Delta/\sigma)_{\text{max}} = 0.001$
$S = 1.08$	$\Delta\rho_{\text{max}} = 0.18 \text{ e \AA}^{-3}$
1761 reflections	$\Delta\rho_{\text{min}} = -0.13 \text{ e \AA}^{-3}$
159 parameters	Extinction correction: <i>SHELXL97</i>
H atoms treated by a mixture of independent and constrained refinement	Extinction coefficient: 0.030 (4)

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$D—H \cdots A$	$D—H$	$H \cdots A$	$D \cdots A$	$D—H \cdots A$
$\text{O2}—\text{H2A} \cdots \text{O1}^i$	1.03 (2)	1.63 (2)	2.657 (2)	174 (2)

Symmetry codes: (i) $-x + 2, -y, -z + 2$.

All C-bound H atoms were positioned geometrically and refined as riding on their carrier atoms, with $\text{C}—\text{H} = 0.93–0.97 \text{ \AA}$ and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{carrier atom})$. The carboxy H atom was located in a difference electron-density map and refined isotropically. The small crystal used in the analysis diffracted rather poorly. The data set was limited to a maximum θ angle of 23.3° , since only very weak diffraction was observed at higher angles.




Figure 2

The crystal packing in (I), showing the strong $\text{O}—\text{H} \cdots \text{O}$ hydrogen bond between facing carboxylic acid groups of molecules related by an inversion center (dashed lines). The asterisk (*) indicates the symmetry operation $(-x, 2 - y, -z)$. H atoms not involved in the interactions have been omitted.

Data collection: *SMART* (Siemens, 1995); cell refinement: *SAINT* (Siemens, 1995); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS86* (Sheldrick, 1985); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *PLATON* (Spek, 2003); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PARST* (Nardelli, 1995).

Thanks are expressed to Arvee Chem Pharma, Mysore, India, for providing the sample. The data collection facility was kindly provided by Dr G. U. Kulkarni at JNC, Jakkur, Bangalore.

References

- Barbui, C., Guaiana, G. & Hotopf, M. (2004). *Pharmacopsychiatry*, **37**, 93–97.
- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bragg, W. H. (1921). *Proc. Phys. Soc.* **34**, 33–50.
- Bragg, W. H. (1922). *J. Chem. Soc.* pp. 2766–2787.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Maizels, M. & McCarberg, B. (2005). *Am. Fam. Physician*, **71**, 483–490.
- Nardelli, M. J. (1995). *J. Appl. Cryst.* **28**, 659.
- Parkin, A., Harte, S. M., Carmichael, D., Currie, S., Drummond, L., Haahr, A., Haggarty, H., Hunter, T., Lamarque, A., Lawton, L., Martin, C., Mathieson, J. E., Mathieson, J. S., McGlone, T., McGregor, J. *et al.* (2005). *Acta Cryst. E* **61**, o2280–o2282.
- Ramesha, A. R. & Roy, A. K. (2001). *Synth. Commun.* **31**, 2419–2422.
- Sheldrick, G. M. (1985). *SHELXS86*. University of Göttingen, Germany.
- Sheldrick, G. M. (1997). *SHELXL97*. University of Göttingen, Germany.
- Siemens (1995). *SMART* and *SAINT*. Version 5.05. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Sim, G. A., Robertson, J. M. & Goodwin, T. H. (1955). *Acta Cryst.* **8**, 157–164.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.